

[From the Proceedings of the Linnean Society of New South Wales,  
1911., Vol. xxxvi., Part 2, June 28th.]

## NOTES ON SOME MALLOPHAGAN GENERIC NAMES.

BY T. HARVEY JOHNSTON, M.A., D.Sc., AND LAUNCELOT  
HARRISON.

A certain amount of confusion has crept into the generic nomenclature of the Insects belonging to the order Mallophaga. Many of the errors have been corrected by Neumann(1906), but there are, however, a few important points in which we differ from the well known French parasitologist. In order that the matters referred to by us might be more easily understood, a very brief sketch of the previous work on certain generic names in this group is necessary.

The difference between "biting lice" and "sucking lice" was first recognised by De Geer(1778), who founded the genus *Ricinus* to include the mandibulate forms, the old name *Pediculus* being retained for the true lice, whose affinities are rather with the Hemiptera than with any other order. Hermann(1804), knowing *Ricinus* to be preoccupied in Botany, proposed *Virmus* as a substitute. Nitzsch, in 1818, published a very important work on parasitic Insecta, including the Mallophaga, but took no notice of previous attempts at classification, beyond placing the earlier generic names as synonyms of his own. He divided the order into two families, to which, however, he did not assign names, these being first given, in 1832, by Burmeister. Each family consisted of two genera, one parasitic on mammals and one on birds. The bird-infesting genera were subdivided by Nitzsch into a number of subgenera as follows:—

Fam. PHILOPTERIDÆ: (1) *Trichodectes*; (2) *Philopterus*(subgenera: *Docophorus*, *Virmus*, *Lipeurus*, *Goniodes*).

Fam. LIOTHEIDÆ: (1) *Gyropus*; (2) *Liotheum*(subgen.: *Colpoccephalum*, *Menopon*, *Trinoton*, *Eureum*, *Lambothrion*, *Physostomum*).

In the case of the two subdivided genera, all species were left in the original genus in each case, but were grouped under the

headings of the various subgenera. These subgenera, though ostensibly retaining that rank, were practically treated as genera by Burmeister(1832), Denny(1842), Giebel(1874), Piaget(1880, 1885), and Taschenberg(1882). It was Kellogg(1896, p.60), who first openly raised the subgenera to genera.

Neumann(1896, pp.56, 60) has pointed out that Hermann had no real justification for erecting *Nirmus*, hence his genus must be ranked as a synonym of *Ricinus*. Moreover, *Nirmus* Nitzsch (1818) could not stand, as the name had already been used by Hermann. Neumann accordingly substituted *Degeeriella* for Nitzsch's genus. He went on to point out that, in the elevation of the subgenera to genera, the original genera had not been retained, and he, therefore, took the first described species under the first named subgenus in each case as the type of the original genus. Thus *Docophorus* was replaced by *Philopterus* (with *P. ocellatus* Scop., as the type), and *Colpocephalum* by *Liotheum* (with *L. zebra* N., as the type). Finally, he considered the question of reinstating *Ricinus*. He has credited Piaget with pointing out that the first species described by De Geer, under his genus as *Ricinus fringilla*, was subsequently included in *Physostomum* N., whereas that author, as well as Giebel and Denny, have, in this matter, simply followed the synonymy given in Nitzsch's work. *Ricinus fringilla* De Geer, was made the type-species of the genus by Neumann, and was said to be probably identical with *Physostomum irascens* N. Nitzsch, Denny, and Giebel have, however, definitely placed *R. fringilla* as a synonym of *Ph. nitidissimum* N., admitting the identity of the two forms. No reasons are advanced by Neumann to show that the latter admission is incorrect. We may therefore conclude that Nitzsch's *Ph. nitidissimum*, rather than his *Ph. irascens*, is the same species as that named by De Geer as *R. fringilla*, and designated as type of *Ricinus* by Neumann. This author is correct in regarding *Physostomum* as a synonym of the latter genus; but it appears to us that the fate of the genera *Liotheum* and *Colpocephalum* is also dependent on the above facts.

Our view of the matter will be seen from the following statement. De Geer founded the genus *Ricinus*, but, as no type had

been designated and all the species had been allotted to later genera, Neumann was at liberty to designate a type from amongst De Geer's original species. *R. fringillæ* having been selected, *Physostomum* N., becomes a synonym of *Ricinus* on account of the former including the type-species of the latter amongst its original species. Nitzsch, however, regarded *Physostomum* as a subgenus, and consistently referred to every species under its generic and not under its subgeneric name. Thus *Ricinus fringillæ* De G., appeared in Nitzsch(1818) as *Liotheum nitidissimum*. Hence *Liotheum* N., includes the designated type of the earlier genus *Ricinus*, and must be ranked as a synonym of the latter. We do not think that it is permissible to fix a type for *Liotheum* as well as for *Ricinus*, seeing that the type of the latter is already included in the former. We have consulted papers by Stiles(1905), and Blanchard(1906), but have not noticed any direct ruling on the point involved. Stiles(1905, p.26), in writing of subdivided genera, states that "if a type is designated, the original generic name always follows the subdivision containing the type." Thus *R. fringillæ* must pass into *Liotheum* before it can reach *Physostomum*. The last-mentioned must be the type-subgenus (in the Nitzschian sense) of *Liotheum*, i.e., of *Ricinus* since it includes the type of the genus. *Colpocephalum* N., would thus maintain its individuality.

Apart from the question of nomenclatural propriety, there can be no doubt as to the expediency of our view, for we would retain the well known genus *Colpocephalum*, which included 137 species up to 1908(Kellogg, 1908), whereas Neumann would reduce it to a synonym of *Liotheum*, a generic name which has remained practically unused since Nitzsch's day. We would place both *Liotheum* N., and *Physostomum* N., as synonyms of *Ricinus*, thus involving a change in the generic names of the twenty-nine species hitherto included under *Physostomum* (vide Kellogg, 1908), a change that is inevitable, as has already been pointed out by Neumann. *Liotheum* Neum., nec Nitzsch, accordingly becomes a synonym of *Colpocephalum*.

Among certain phlopterid genera, an extraordinary confusion has arisen, the various stages of which are as follows. In 1835,

Dufour described a parasite, *Phlopterus brevis*, from an albatross, this species being made the type of the genus *Docophoroides* Denny(MSS.) by Giglioli in 1864, a valid generic diagnosis being given. Nitzsch had named the same insect *Lipeurus taurus*, but since his description was not published until 1866 (p.385), it cannot claim precedence. Subsequent writers apparently recognise these facts in their synonymy, but Denny's genus has not received due recognition. In 1866, Rudow founded *Trabeculus* for a closely allied parasite, *Tr. schillingi*, from a petrel; and, four years later, ignoring his own prior action, created a new genus, *Oncophorus* (nec *Oncophora* Dies., a nematode genus) with the same type. In 1882, Taschenberg erected *Eurymetopus* with *Lipeurus taurus* N., as type, recognised Rudow's *Oncophorus schillingi* as congeneric, after having examined the type-specimen of the latter, and placed this species under his own genus. Piaget, in 1880, added several species to *Oncophorus* Rudow, dividing them into two groups, the *Docophoroides* and the *Nirmoides*; but he was obviously not too sure of his ground, since, though he included Rudow's species in his descriptions, he did not include it in his key to the genus. Taschenberg having removed *O. schillingi* from the genus *Oncophorus* into his *Eurymetopus*, Piaget in 1885 (p.35) claimed the genus as "*Oncophorus*, m." Kellogg(1896, p.68) has indicated some of these facts, but has not offered any comment, while Neumann appears to have overlooked them, and to have accepted Piaget's genus as being identical with Rudow's *Trabeculus*, which name he allowed to stand. He gave a new name, *Taschenbergius*, in place of *Eurymetopus*, a name already preoccupied as a genus of Coleoptera by Schönherr in 1840. From the foregoing it is obvious that *Docophoroides* Denny, stands, with *Trabeculus* Rud., *Oncophorus* Rud., *Eurymetopus* Tasch., and *Taschenbergius* Neum., as synonyms, for the forms from petrels: while a new generic name is required for the species from *Rallidæ*, etc., grouped by Piaget and later writers under *Oncophorus* (*O. schillingi* and allied *Docophoroid* forms being excluded). For this group of species, we propose the name *Rallicola*, which differs etymologically from *Rallicula* Schleg., 1871, in ornithology. *O. attenuatus* N., has been selected by us as the type-species.

Neumann has already replaced *Nitzschia* Denny, 1842 (*nec* v. Baer, 1827) by *Dennyus*, and *Piagetia* Picaglia, 1885, (*nec* Ritsenia, 1874) by *Piagetiella*. Picaglia's paper is not available to us, and his genus has not been taken into account by Kellogg, beyond the mere mention of it in his bibliography (1896, p.36). The species are listed under *Menopon* in the latter's list (Kellogg, 1908). We are, therefore, not able to give any opinion as to the value of the genus. A subdivision of the unwieldy *Menopon* is, however, desirable. *Eureum* N., is included by Neumann in his list of genera, though Kellogg (1899, p.133) has shown it to be based on immature forms of a *Menopon*. The latter author has studied *E. malleus* N., (= *M. malleus*), which, being the better known of Nitzsch's two species, may be taken as the type.

In a later publication, Neumann (1909, p.9, footnote) has changed *Ornithobius* Denny, to *Ornithonomus*, as he considers the former name to be preoccupied by *Ornithobia* Meigen, 1832, in Diptera. In the recommendations under Art.36 of the International Code, as given by Stiles (1905, p.47), and by Blanchard (1906, p.32), this is not allowed to be sufficient grounds for the rejection of a name. We have accordingly placed *Ornithonomus* Neum., as a synonym of *Ornithobius* Denny, in the list of genera appended. Piaget (1880, p.378) has already shown that *Metapeuron* Rudow, (type *M. punctatum* Rud., = *O. cygni* L.) is a synonym of this genus.

The following genera, proposed prior to Neumann's paper, have been omitted by him, viz., *Nesiotinus* Kellogg, 1903; *Phloceanus* Kell., 1903; *Ornicholax* Carriker, 1903; *Kelloggia* Carriker, 1903; *Heterodoxus* Le Souëf & Bullen, 1902; *Latumcephalum* Le Souëf, 1902.

In conclusion, we append a list of the twenty-eight genera at present existing, indicating the types where our available literature has permitted. Such of these types as are marked with an asterisk(\*), have been designated as such by ourselves, and, for the most part, have been selected from the available species on account of their occurrence on common domestic animals, or on other hosts which are fairly readily obtainable.

## A. Suborder ISCHNOCERA Kellogg.

F a m. *Trichodectidæ*:—

1. *Trichodectes* N., 1818.—\**T. canis* Degeer (syn. *T. latus* N.).

F a m. *Philopteridæ*:—

2. *Philopterus* N., 1818 (syn. *Docophorus* N., 1818)—*P. ocellatus* Scop.  
 3. *Vesiotinus* Kell., 1903.—*V. demersus* Kellogg.  
 4. *Kelloggia* Carriker, 1903.—*K. brevipes* Carr.  
 5. *Ornicholax* Carriker, 1903.—*O. robustus* Carr.  
 6. *Akidoproctus* Piaget, 1878.—*A. marginatus* P.  
 7. *Giebelia* Kellogg, 1896.—*G. mirabilis* K.  
 8. *Degeeriella* Neum., 1906 (syn. *Nirmus* N., 1818, nec Herm., 1804).—\**D. discocephalus* N.  
 9. *Docophoroides* Denny in Giglioli, 1864 (syns. *Trabeculus* Rud., 1866; *Oncophorus* Rud., 1870, nec Piaget, 1885; *Eurymetopus* Tasch., 1882, *Taschenbergius* Neum., 1906).—*D. brevis* Dufour.  
 10. *Goniodes* N., 1818.—\**G. pavonis* L. (syn. *G. falcicornis* N.).  
 11. *Goniocotes* Burm., 1835.—\**G. gallinæ* Deg. (syn. *G. hologaster* N.).  
 12. *Ornithobius* Denny, 1842 (syn. *Metapeuron* Rud., 1870; *Ornithonomus* Neum., 1909).—*O. cygni* Linn.  
 13. *Bothriometopus* Tasch., 1882.—*B. macrocnemis* N.  
 14. *Philoceanus* Kellogg, 1903.—*P. becki* K.  
 15. *Lipecurus* N., 1818.—\**L. caponis* Linn. (syn. *L. variabilis* N.).  
 16. *Rallicola* nobis, 1911 (syn. *Oncophorus*, Piag., 1885; nec Rudow, 1870).—\**R. attenuatus* N.

## B. Suborder AMBLYCERA Kellogg.

F a m. *Gyropidæ*.

17. *Gyropus* N., 1818.—\**G. ovalis* N.

F a m. *Ricinidæ*.†

18. *Ricinus* Degeer, 1778 (syns., *Nirmus*, Herm., 1804, nec N., 1818; *Liotheum* N., 1818, nec Neum., 1906; *Physostomum* N., 1818).—*R. fringillæ* Deg.
19. *Colpocephalum* N., 1818 (syn. *Liotheum* Neum., 1906, nec N., 1818).—*C. zebra* N.
20. *Menopon* N., 1818 (syn. *Eureum* N., 1818).—\**M. gallinæ* L. (syns. *M. trigonocephalum* Olfers; *M. pallidum* N.).
21. *Trinoton* N., 1818.—\**T. anseris* Sulzer (syn. *T. conspurcatum* N.).
22. *Læmbothrion* N., 1818.—\**L. maximum* Scop. (syn. *L. giganteum* N.).
23. *Boopia* Piaget, 1880.—*B. tarsata* Piag.
24. *Dennyus* Neum., 1906 (syn. *Nitzschia* Denny, 1842, nec v. Baer, 1827).—*D. burmeisteri* Denny (syn. *N. pulicaris* N., in Giebel, 1874).
25. *Ancistrona* Westwood, 1874.—*A. procellariæ* Westw.
26. *Piagetiella* Neum., 1906, (syn. *Piagetia* Picaglia 1885, nec Ritsenia).
27. *Heterodoxus* Le Souëf & Bullen, 1902.—*H. macropus* Le Souëf & Bullen.
28. *Latumcephalum* Le Souëf, 1902.—*L. macropus* Le Souëf.

## BIBLIOGRAPHY.

- 1906, BLANCHARD—"Règles internat.d.l.Nomencl. Zool. etc." Bull. Soc. Zool. France, xxxi., 1906, pp.15-35.
- 1832, BURMEISTER—Handb. d. Entomol. 1832.
- 1778, DE GEER—Mémoires pour servir à l'histoire des Insectes, vii., p. 69.
- 1842, DENNY—Monogr. Anoplurorum Britannia, London, 1842.
- 1835, DUFOUR—Ann. Soc. Ent. France, iv., 1835.
- 1874, GIEBEL—"Insecta Epizoa etc.," Liepzig, 1874.
- 1864, GIGLIOLI—Q.J.M.S. iv., 1864.
- 1804, HERMANN—Mémoire aptérologique, xii., 1804, p.12.

† Called *Liotheinæ* in Neumann(1906, pp.58, 60). *Ricinidæ* is the more correct designation, as the name refers to a family whose type-genus is *Ricinus*.

- 1896, KELLOGG—"New Mallophaga, i." Proc. Calif. Acad. Sci.(2), vi., 1896, and in Contributions to Biology from the Hopkins Seaside Lab., iv., 1896.
- 1899, ————"New Mallophaga, iii." Proc. Calif. Acad. Sci., Occas. Papers, vi., 1899.
- 1908, ————"Mallophaga," in Wyttsmann's "Genera Insectorum." Fasc. 66.
- 1906, NEUMANN—Bull. Soc. Zool. France, xxxi., 1906, pp. 54, *sqg.*
- 1909, ————"Parasites et Maladies parasitaires des Oiseaux domestiques." Paris, 1909.
- 1818, NITZSCH—in Germar's Magazin d. Entom., iii., 1818, pp. 261, *sqg.*
- 1866, ————Zeitschr. f. ges. Naturwiss. xxviii.
- 1874, ————in Giebel's "Insecta Epizoa," 1874.
- 1880, PIAGET—"Les Pediculines, Essai Monographique." Leiden, 1880.
- 1885, ———— " " Supplement. Leiden, 1885.
- 1884, PICAGLIA—Atti d. Soc. d. Nat. d. Modena, (3), ii., 1884, p. 103.
- 1866, RUDOW—Zeitschr. f. ges. Naturwiss. xxvii., 1866, p. 466.
- 1870, ———— " " xxxv., 1870, p. 138.
- 1905, STILES—Bull. 24, Hyg. Lab. Publ. Health & Mar. Hosp. Service, U.S.A.
- 1882, TASCHENBERG—"Die Mallophagen etc." Nova Acta, Halle, xlv., 1, 1882.

*Postscript, added 15th July, 1910.*—After this paper was read, we came upon a reference in Kellogg(1896, p.164) which gives the generic characters of *Piagetia*, *P. ragazzii* Picaglia, being the type. This parasite is certainly congeneric with *Menoion titan* Piaget, and *Tetrophthalmus chilensis* Grosse(Zeit. f. wiss. Zool., xlii., 1885, p.534). As Grosse's genus *Tetrophthalmus* antedates Picaglia's *Piagetia*, which name is, moreover, already preoccupied, as Neumann has pointed out, Grosse's genus must stand (with *T. chilensis* as type), and *Piagetia* Picaglia, as well as *Piagetiella* Neumann, must rank as synonyms. We refrain at present from commenting upon the individuality of *T. chilensis*, and the various other species and subspecies of the *titan*-group (*T. titan*, Piag., *T. ragazzii* Pic., *T. consanguineus* Piag., *T. linearis* Kell., and *T. impar* Kell.).

According to the ruling of the International Commission, Grosse's genus *Tetrophthalmus* is not invalidated by *Tetrophthalma* Less., 1833(Coleoptera), and *Tetraophthalmus* de Haan, 1834 (Coleoptera).



*[From the Proceedings of the Linnean Society of New South Wales,  
1911., Vol. xxxvi., Part 2, June 28th.]*

---

NOTES ON SOME MALLOPHAGAN GENERIC NAMES.

BY T. HARVEY JOHNSTON, M.A., D.Sc., AND LAUNCELOT  
HARRISON.

[Issued 20th November, 1911.]