

J. Raptor Res. 40(4):301–302

© 2006 The Raptor Research Foundation, Inc.

LICE OF CHILEAN OWLS: A FIRST DESCRIPTION

DANIEL GONZÁLEZ-ACUÑA AND RODRIGO MUÑOZ C.

Facultad de Medicina Veterinaria, Universidad de Concepción, Casilla 537, Chillán, Chile

ARMANDO CICCHINO

Castelli 4117, 7600 Mar del Plata, Argentina

RICARDO A. FIGUEROA R.

Instituto de Ecología y Evolución, Escuela de Graduados, Facultad de Ciencias, Universidad Austral de Chile, Casilla 567, Valdivia, Chile

and

Estudios para la Conservación y Manejo de la Vida Silvestre Consultores, Blanco Encalada 350, Chillán, Chile

KEY WORDS: *Kurodaia*; *Strigiphilus*; *Chile*; lice; owls.

Lice (Phthiraptera) are obligate ectoparasitic insects, carrying out their entire life cycle on the same mammal or bird host. Under particular conditions, lice may have negative effects on their hosts such as decreasing thermoregulatory capacity, reducing nestlings' body mass (which may negatively affect viability and productivity), influencing sexual selection, or transmitting pathogenic endoparasites (Clayton 1990, Moller 1990, Hunter et al. 1994). Lice often multiply when their hosts are stressed, debilitated, or sick (Krone and Cooper 2002). This may reach major importance in terms of conservation and management of threatened mammals and bird species (e.g., Smith and Belthoff 2001). On the other hand, because most genera of the Phthiraptera are restricted to particular taxa and because some louse species parasitize only one host species or a number of subspecies (Clayton 1990, Hahn et al. 2000), they can be useful in understanding ecological, coevolutionary, or phylogenetic relationships (Marshall 1981, Hafner et al. 1994, Hahn et al. 2000, Clayton and Drown 2001). In Chile, diversity and taxonomy of lice associated with wild bird hosts have been only recently studied (González-Acuña et al. 2003a, 2003b, 2004, 2005, San Martín et al. 2005). Here, we report seven lice species parasitizing Chilean owls. In part, these represent new records for Chile.

METHODS

We analyzed a total of 95 birds of six owl species: Barn Owl (*Tyto alba*, $N = 21$), Magellanic Horned Owl (*Bubo magellanicus*, $N = 20$), Austral Pygmy Owl (*Glaucidium nanum*, $N = 22$), Short-eared Owl (*Asio flammeus*, $N = 14$), Burrowing Owl (*Athene cunicularia*, $N = 16$) and Rufous-legged Owl (*Strix rufipes*, $N = 3$). Lice were obtained from several birds held in wildlife rehabilitation centers in central and southern Chile ($N = 26$), from carcasses found as roadkills ($N = 24$), and also from the bird collection housed in the Chilean National Museum of Natural History in Santiago

($N = 45$). In order to remove all louse specimens, each bird or carcass was handled on a clean white surface and the entire plumage was profusely and systematically surveyed (Pérez et al. 1996). All visible lice were removed from the host and brought into the laboratory for examination. The collected material was fixed in 70% filtered ethanol solution, and then slide-mounted for identification following the technique published by Palma (1978) and Price et al. (2003). This material was deposited in the collection of the Faculty of Veterinary Medicine of the University of Concepción, Chile.

RESULTS AND DISCUSSION

A total of 130 lice of seven species were isolated from all six sampled owl species. Two louse species were isolated from Barn Owls and one each from Short-eared Owls, Burrowing Owls, Magellanic Horned Owls, Austral Pygmy Owls, and Rufous-legged Owls (Table 1). Although *Kurodaia subpachygaster* was previously reported from Barn Owls (Price and Beer 1963), *Strigiphilus symii* on *S. rufipes* (Clayton 1990), and *S. chilensis* on *B. virginianus nacurutu* (renamed *B. magellanicus*, Clayton and Price 1984), all the remaining louse species we found represented new records for Chile. Because most of the owl specimens were obtained from museum and rehabilitation centers, our results probably do not reflect the natural condition, but, nonetheless, are useful in identifying louse species and their host specificity.

PIOJOS DE BÚHOS CHILENOS: UNA PRIMERA DESCRIPCIÓN

RESUMEN.—Examinamos 95 especímenes de búhos obtenidos de centros de rehabilitación de fauna silvestre, aves muertas en carreteras y colecciones de museo. Se analizaron un total de 20 individuos de la especie *Tyto alba*, 20 de *Bubo magellanicus*, 22 de *Glaucidium nanum*, 14 de *Asio flammeus*, 16 de *Athene cunicularia* y 3 de *Strix rufipes*. Dos especies de piojos fueron aisladas de *T. alba* (*Kurodaia subpachygaster*, *Strigiphilus aitkeni*) y una especie de *B. magellanicus* (*S. chilensis*), *G. nanum* (*S. microgenitalis*), *A. flam-*

¹ Email address: danigon@udec.cl

Table 1. Louse species collected from Chilean Owls.

OWL SPECIES	LOUSE SPECIES	ADULT		NYMPH (N)	SEX RATIO (M/F)
		MALE (N)	FEMALE (N)		
<i>Strix rufipes</i>	<i>Strigiphilus syrni</i>	—	7	1	—
<i>Athene cucularia</i>	<i>Strigiphilus speotyti</i>	2	2	—	1:1
<i>Glaucidium nanum</i>	<i>Strigiphilus microgenitalis</i>	1	1	6	1:1
<i>Bubo magellanicus</i>	<i>Strigiphilus chilensis</i>	18	22	10	1:1.2
<i>Asio flammeus</i>	<i>Strigiphilus cursor</i>	5	9	8	1:1.8
<i>Tyto alba</i>	<i>Kurodaia subpachygaster</i>	1	2	—	1:2
	<i>Strigiphilus (Tytoniella) aitheni</i>	11	16	—	1:1.4

meus (*S. cursor*), *A. cucularia* (*S. speotyti*) y *S. rufipes* (*S. syrni*). Las especies *S. speotyti*, *S. microgenitalis*, *S. cursor*, y *S. aitheni* representan nuevos registros para Chile.

[Traducción del equipo editorial]

ACKNOWLEDGMENTS

We thank Eduardo Pavéz, Sergio Alvarado, and Benito González, who kindly permitted access to wildlife rehabilitation centers. J.C. Torres-Mura kindly permitted access to the bird collection in the National Museum of Natural History in Santiago. R.A. Figueroa was supported by a Doctoral grant from Comisión Nacional de Investigación Científica y Tecnológica (CONICYT) and Escuela de Graduados, Facultad de Ciencias, Universidad Austral.

LITERATURE CITED

- CLAYTON, D.H. 1990. Mate choice in experimentally parasitized crock doves: lousy males lose. *Am. Zool.* 30:251–262.
- AND D.M. DROWN. 2001. Critical evaluation of five methods for quantifying chewing lice (Insecta: Phthiraptera). *J. Parasitol.* 87:1291–1300.
- AND R.D. PRICE. 1984. Taxonomy of the *Strigiphilus cursitans* Group (Ischnocera: Philopteridae) parasites of owls (Strigiformes). *Ann. Entomol. Soc. Am.* 77:340–363.
- GONZÁLEZ-ACUÑA, D., A. DAUGSCHIES, K. POLMEYER, L. RUBILAR-CONTRERAS, O. SKEWES-RAMM, AND E. MEY. 2003a. Ectoparasitos de la perdiz chilena (*Nothoprocta perdicaria*) en la provincia de Ñuble, Chile. *Parasitol. Latinoam.* 58:75–77.
- , —, —, —, —, —, AND M. CASANUEVA. 2003b. Ectoparasitos de la codorniz (*Callipepla californica*) en la provincia de Ñuble, Chile y su correlación con el sexo, edad y hábitat de captura. *Lundiana* 4:129–134.
- , —, —, L. RUBILAR-CONTRERAS, K. POLMEYER, O. SKEWES-RAMM, AND E. MEY. 2004. Fauna parasitaria de la tórtola común (*Zenaida auriculata*, de Murs 1847) (Columbiformes: Columbidae) en Ñuble, Chile. *Parasitol. Latinoam.* 59:37–41.
- , O. SKEWES-RAMM, C. CANDIA, R. PALMA, AND L. MORENO. 2005. Estudio del parasitismo gastrointestinal y externo del Caiquén *Chloephaga picta* Gmelin, 1789 (Aves, Anatidae) en la Región de Magallanes, Chile. *Parasitol. Latinoam.* 60:86–89.

HAFNER, M.S., D.S. PHILIP, F.X. VILLABLANCA, T.A. SPRADLING, J.W. DEMASTES, AND S.A. NADLER. 1994. Disparate rates of molecular evolution in cospeciating hosts and parasites. *Science* 265:1087–1090.

HAHN, D.C., R.D. PRICE, AND P.C. OSENTON. 2000. Use of lice to identify cowbird hosts. *Auk* 117:943–951.

HUNTER, E., R.J. GUTIÉRREZ, A.B. FRANKLIN, AND D. OLSON. 1994. Ectoparasites of the Spotted Owl. *J. Raptor Res.* 28:232–235.

KRONE, O. AND J.E. COOPER. 2002. Parasitic diseases. Pages 105–120 in J.E. Cooper [Ed.], *Birds of prey: health and diseases*. Blackwell Science, Oxford, U.K.

MARSHALL, A.G. 1981. The ecology of ectoparasitic insects. Academic Press, New York, NY U.S.A.

MOLLER, A.P. 1990. Effects of parasitism by a haematophagous mite on reproduction in the Barn Swallow. *Ecology* 71:2345–2357.

PALMA, R. 1978. Slide-mounting of lice: a detailed description of the Canada balsam technique. *N. Z. Entomol.* 6:432–436.

PÉREZ, J.M., I. RUIZ-MARTÍNEZ, AND J.E. COOPER. 1996. Occurrence of chewing lice on Spanish raptors. *Ardeola* 43:129–138.

PRICE, R. AND J.R. BEER. 1963. The *Kurodaia* (Mallophaga: Menoponidae) parasitic on the *Strigiformes*, with a key to the species of the genus. *Ann. Entomol. Soc. Am.* 55:849–857.

PRICE, R.D., R.A. HELLENTHAL, R.L. PALMA, K.P. JOHNSON, AND D.H. CLAYTON. 2003. The chewing lice: world checklist and biological overview. Illinois Natural History Survey Special Publication 24, Champaign, IL, U.S.A.

SAN MARTÍN, J.C. BREVIS, L. RUBILAR, R. SCHMASCHKE, A. DAUGSCHIES, AND D. GONZÁLEZ-ACUÑA. 2005. Ectoparasitismo en tique común *Milvago chimango chimango* (Viellot, 1816) (Aves, Falconidae) en la zona de Ñuble, Chile. *Lundiana* 6:49–55.

SMITH, B.W. AND J.R. BELTHOFF. 2001. Identification of Ectoparasites on Burrowing Owls in southwestern Idaho. *J. Raptor Res.* 35:159–60.

Received 28 January 2006; accepted 16 October 2006