

Revised by
K. C. Emerson
15 November 1971

MALLOPHAGA GENERA AND SPECIES
FOUND ON MAMMALS

BOOPIDAE

Boopia

- bettongia LeSouëf, 1902
(phanerocerata Harrison and Johnston, 1916)
- biseriata Keler, 1971
- doriana Keler, 1971
- dubia Werneck and Thompson, 1940
- grandis Piaget, 1885
(peregrina Mjöberg, 1910)
- minuta LeSouëf, 1902
- mjöbergi Werneck and Thompson, 1940
- notafusca LeSouëf, 1902
- tarsata Piaget, 1880
- uncinata Harrison and Johnston, 1916

Heterodoxus

- alatus Keler, 1971
- ampullatus Keler, 1971
- ancoratus Keler, 1971
- calabyi Keler, 1971
- longitarsus (Piaget, 1880)
- maai Emerson, 1962
- macropus LeSouëf and Bullen, 1902
- mitratus Keler, 1971
- octoseriatus Keler, 1971
- pygidialis (Mjöberg, 1919)
- quadriseriatus Keler, 1971
- spiniger (Enderlein, 1909) *
(armiferus Paine, 1912)
- ualabati Plomley, 1940

Latumcephalum

- lesouëfi Harrison and Johnston, 1916
- macropus LeSouëf, 1902

Macropophila

- biarcuata Keler, 1971
- breviarcuata Keler, 1971
- clayae Keler, 1971
- forcipata Mjöberg, 1919

Paraboopia
flava Werneck and Thompson, 1940

Paraheterodoxus
calcaratus Keler, 1971
erinaceus Keler, 1971
insignis Harrison and Johnston, 1916

Phacogalia
brevispinosa Harrison and Johnston, 1916
spinosa Harrison and Johnston, 1916

GYROPIDAE

Aotiella
aotophilus Ewing, 1924 *

Gliricola
aequatorialis Werneck, 1948
arboricola Mendez, 1969 *
brasiliensis Werneck, 1934
calcaratus Werneck, 1937
capromydis armatus Werneck, 1944 *
capromydis capromydis Werneck, 1944 *
columbanus Werneck, 1944
cubanus Werneck, 1944 *
decurtatus decurtatus (Neumann, 1912)
decurtatus fonsecai Werneck, 1933
decurtatus maculatus Werneck, 1942
decurtatus marajoensis Werneck, 1942 *
decurtatus paraensis Werneck, 1942
distinctus Ewing, 1924
echimydis Werneck, 1933
ewingi Werneck, 1944 *
humilis Werneck, 1951
lindolphoi Werneck, 1942
longicollis (Burmeister, 1838)
mesomydis Werneck, 1938
mirandi Werneck, 1935
o'mahonyi Werneck, 1951 *
palladius Keler, 1943
panamensis Werneck, 1944 *
pintoi Werneck, 1935
porcelli (Schrank, 1781) *

* Known from North America (North of Colombia)

Gliricola (Continued)
(saviae Schrank, 1803)
(bifurcatus Olfers, 1816)
(gracilis Nitzsch, 1818)
(porcelli perfoliatus Neumann, 1912)
(bicaudatus Paine, 1912)
(mexicanus Zavaleta, 1946)
quadrisetosus (Ewing, 1924)
sylvantica Mendez, 1969 *
vogelsangi Werneck, 1951

Gyropus
cercomydis Werneck, 1942
cruzi (Werneck, 1933)
diplomys Mendez, 1967 *
emersoni Mendez, 1969 *
freitasi Werneck, 1942
lenti distinctus Werneck, 1948
lenti lenti Werneck, 1936
limai Werneck, 1948
lineatus Neumann, 1912
longus Neumann, 1912
martini iheringi Werneck, 1948
martini martini (Werneck, 1934)
martini matthaeensis Werneck, 1948
mesoamericanus Mendez, 1969 *
ovalis Burmeister, 1838 *
 (turbinatus Piaget, 1880)
 (recifensis Torres, 1938)
 (mexicanus Zavaleta, 1945)
parasetosus Werneck, 1935
parvus (Ewing, 1924)
ribeiroi Werneck, 1935
scalaris Werneck, 1942
setifer (Ewing, 1924) *
setosus Neumann, 1912
thompsoni Werneck, 1935
travassosi Werneck, 1948

Macrogyropus
amplexans amplexans (Neumann, 1912) *
 (ratti Larranaga, 1924)
amplexans longisetis Werneck, 1948
costalimai (Werneck, 1931) *

* Known from North America (North of Colombia)

Macrogyropus (Continued)
dicotylis (Macalister, 1869) *
 (dentatus Ewing, 1924)
heteronychus (Ewing, 1924)

Monothoracius
almeidai Werneck, 1937
penidoi Werneck, 1934

Phtheiropoios
ewingi (Werneck, 1936)
 (forficulatus Ewing, 1924)
 (grypophallus Werneck, 1939)
forficulatus (Neumann, 1912)
lagotis (Gervais, 1849)
 (alpinus Kellogg and Nakayama, 1914)
latipollicaris (Ewing, 1924)
 (gracilipes Ewing, 1924)
 - (latipollicaris Werneck, 1936)
nematophallus (Werneck, 1935)
pearsoni Werneck, 1948
pollicaris (Ewing, 1924)
wetmorei (Ewing, 1924)

Pitrufoquia
coypus Marelli, 1932 *
 (mollis Marelli, 1933)

Protogyropus
normalis Ewing, 1924

HAEMATOMYZIDAE

Haematomyzus
elephantis Piaget, 1869
 (proboscideus Piaget, 1880)
 (paradoxus Lahille, 1906)
 (sumatranus Fahrenholz, 1910)
hopkinsi Clay, 1963

PHILOPTERIDAE

Trichophlopterus

* Known from North America (North of Colombia)

Trichophlopterus (Continued)
babakotophilus Stobbe, 1913
(ferrisi Eichler, 1942)
(stobbei Mjöberg, 1919)

TRICHODECTIDAE

Bovicola
adenota Bedford, 1936
alpina Keler, 1942
(rubicaurae Werneck, 1944)
asini (Eichler, 1953)
aspilopyga Werneck, 1956
breviceps (Rudow, 1866)
(inaequalemaculatus Piaget, 1885)
(aucheniae - lamae Torreggiani, 1910)
bovis (Linnaeus, 1758) *
(tauri Linnaeus, 1761)
(scalaris Nitzsch, 1818)
caprae (Gurlt, 1843) *
(climax Nitzsch, 1861)
(climacium Giebel, 1861)
(solidus Rudow, 1866)
(mambricus Rudow, 1866)
(caprae Packard, 1870)
(climax truncata Piaget, 1880)
(peregrinus Taschenberg, 1882)
connectens (Hopkins, 1943)
conconvifrons (Hopkins, 1960) *
crassipes (Rudow, 1866) *
(penicillatus Piaget, 1880)
(hermsi Kellogg and Nakayama, 1915)
(ibicis Gaschen, 1951)
dimorpha Bedford, 1939
equi (Denny, 1842) *
(pilosus Giebel, 1874)
(parumpilosum Piaget, 1880)
(parumpilosum tarsatus Piaget, 1880)
(pubescens Macqueen, 1905)
(vestitus Railliet, 1895)
hemitrangi (Cummings, 1916)
hilli Bedford, 1934
jellisoni Emerson, 1962 *
limbata (Gervais, 1844) *

* Known from North America (North of Colombia)

Bovicola (Continued)
(climax major Piaget, 1885)
(madagascariensis Mjöberg, 1910)
(painei Kellogg and Nakayama, 1914)
(sachtlebeni Keler, 1937)
(wernecki Keler, 1937)
longicornis (Nitzsch, 1818) *
(similis Denny, 1843)
(americanus Jellison, 1935)
neglecta Keler, 1942
ocellata (Piaget, 1880)
oreamnidis (Hopkins, 1960) *
ovis (Schrank, 1781) *
(ovisarietis Schrank, 1803)
(sphaerocephalus Olfers, 1816)
(sphaerocephalus Nitzsch, 1818)
pelea Bedford, 1934
sedecimdecembrii Eichler, 1946 *
tarandi (Mjöberg, 1910) *
thompsoni Bedford, 1936
tibialis (Piaget, 1880)
(longicornis Denny, 1842)
(tibialis alba Piaget, 1880)
(punctum Piaget, 1885)
trampel (Eicherl, 1954)
zuluensis Werneck, 1950

Cebidicola

armatus (Neumann, 1913) *
extrarius Werneck, 1950
semiarmatus (Neumann, 1913) *

Damalinia

appendiculata (Piaget, 1880)
baxi Hopkins, 1947
chlorleyi (Hopkins, 1941)
crenelata (Piaget, 1880)
forficula forficula (Piaget, 1880)
forficula muntiacus (Seguy, 1948)
(siamensis Werneck, 1950)
harrisoni (Cummings, 1916)
hendrickxi Hopkins, 1947
hopkinsi Bedford, 1936
martinaglia (Bedford, 1936)
meyeri (Taschenberg, 1882)

* Known from North America (North of Colombia)

Damalinia (Continued)

(hamadryas Fahrenholz, 1919)
natalensis Emerson, 1963
neotheileri Emerson and Price, 1971
ornata Werneck, 1957
semitheileri Emerson and Price, 1971
theileri Bedford, 1928
traguli Werneck, 1950

Dasyonyx (Dasyonyx)

bedfordi Werneck, 1935
dendrohyracis (Ferris, 1930)
guineensis Werneck, 1941
hopkinsi Werneck, 1941
minor Bedford, 1939
ocuiatus (Bedford, 1928)
ovalis Bedford, 1932
(windhuki Bedford, 1936)
smallwoodae Emerson and Price, 1969
validus ugandensis Werneck, 1941
validus validus Bedford, 1932
(lindfieldi Ferris, 1930)

Dasyonyx (Neodasyonyx)

capensis Emerson, 1964
discanthus (Ehrenberg, 1828)
nairobiensis Bedford, 1936
ruficeps Emerson, 1964
transvaalensis Bedford, 1932
waterbergensis Bedford, 1932

Eurytrichodectes

machadoi Werneck, 1958
paradoxus Stobbe, 1913

Eutrichophilus

cercolabes Mjöberg, 1910
(australis Ewing, 1936)
comitans Werneck, 1950
cordiceps Mjöberg, 1910
exiguus Werneck, 1950
guyanensis Werneck, 1950
lobatus Ewing, 1936
maximus Bedford, 1939 *

*Known from North America (North of Colombia)

Eutrichophilus (Continued)

mexicanus (Rudow, 1866) *
(coëndu Stobbe, 1913)
minor Mjöberg, 1910
moojeni Werneck, 1945
setosus (Giebel, 1861) *

Felicola

acutirostris (Stobbe, 1913)
bedfordi Hopkins, 1942
caffra (Bedford, 1919)
calogaleus (Bedford, 1928)
cynictis (Bedford, 1928)
felis (Werneck, 1934) *
genettae (Fresca, 1924) (nec Bedford, 1919)
hercynianus Koler, 1957
inaequalis (Piaget, 1880)
(rammei Stobbe, 1913)
intermedius hyaenae Hopkins, 1960
intermedius intermedius (Bedford, 1932)
jucci (Conci, 1942)
macrurus Werneck, 1948
minimus Werneck, 1948
pygidialis Werneck, 1948
rahmi Emerson and Stojanovich, 1966
robertsi Hopkins, 1944
rohani Werneck, 1956 *
setosus Bedford, 1932
siamensis Emerson, 1964
spenceri Hopkins, 1960 *
subrostratus (Burmeister, 1838) *
(rostratus Bedford, 1932)
zeylonicus Bedford, 1936

Geomydoecus

alcorni Price and Emerson, 1971 *
alleni Price and Emerson, 1971 *
bulleri Price and Emerson, 1971 *
californicus (Chapman, 1897) *
chapini Werneck, 1945 *
chiapensis Price and Emerson, 1971 *
copei Werneck, 1945
coronadoi Barrera, 1961 *
costaricensis Price and Emerson, 1971 *

* Known from North America (North of Colombia)

Geomydoecus (Continued)

dakotensis Price and Emerson, 1971 *
dariensis Price and Emerson, 1971 *
duchesnensis Price and Emerson, 1971 *
ewingi Price and Emerson, 1971 *
expansus (Duges, 1902) *
fulvescens Price and Emerson, 1971 *
geomydis geomydis (Osborn, 1891) *
geomydis subgeomydis Price and Emerson, 1971 *
jonesi Price and Emerson, 1971 *
illinoensis Price and Emerson, 1971 *
mcgregori Price and Emerson, 1971 *
merriami Price and Emerson, 1971 *
mexicanus Price and Emerson, 1971 *
minor Werneck, 1950 *
neocopei Price and Emerson, 1971 *
oklahomensis Price and Emerson, 1971 *
oregonus idahoensis Price and Emerson, 1971 *
oregonus oregonus Price and Emerson, 1971 *
panamensis Price and Emerson, 1971 *
perotensis irolonis Price and Emerson, 1971 *
perotensis perotensis Price and Emerson, 1971 *
polydentatus Price and Emerson, 1971 *
quadridentatus Price and Emerson, 1971 *
scleritus (McGregor, 1917) *
subcalifornicus Price and Emerson, 1971 *
texanus Ewing, 1936 *
thomomyus (McGregor, 1917) *
tolucae Price and Emerson, 1971 *
trichopi Price and Emerson, 1971 *
truncatus Werneck, 1950
umbrini Price and Emerson, 1971 *
veracruzensis Price and Emerson, 1971 *
wardi Price and Emerson, 1971 *
wernecki Price and Emerson, 1971 *
yucatanensis Price and Emerson, 1971 *

Lorisicola

mjöbergi (Stobbe, 1913)
(brachycephalus Ewing, 1930)

Lutridia

exilis (Giebel, 1861) *
lutrae (Werneck, 1937)
matschiei (Stobbe, 1913)

* Known from North America (North of Colombia)

Lymcon

cummingsi Eichler, 1943 *
gastrodes (Cummings, 1916) *

Neofelicola

aspidorhynchus Werneck, 1948
bengalensis Werneck, 1948
philippiensis Emerson, 1965
sumatrensis Werneck, 1948

Neotrichodectes

arizonae Werneck, 1948 *
(spatulatus Cook, 1953)
chilensis Werneck, 1948
interruptofasciatus (Kellogg and Ferris, 1915) *
mephitidis (Packard, 1872) *
(castoris Osborn, 1896)
(monticulus McGregor, 1917)
minutus (Paine, 1912) *
osborni Keier, 1944 *
(mephitidis Osborn, 1896)
pallidus (Piaget, 1880) *
(nasuatis Osborn, 1902)
thoracicus (Osborn, 1902) *
wolffhugeli (Werneck, 1936)

Parafelicola

acuticeps (Neumann, 1902)
(genetta Bedford, 1919)
africanus Emerson and Price, 1966
lenicornis Werneck, 1948
neoafricanus Emerson and Price, 1966
viverriculae (Stobbe, 1913)
wernecki (Hopkins, 1941)

Procavicola (Condylocephalus)

bedfordi bedfordi Werneck, 1941
bedfordi dissimilis Werneck, 1941
hopkinsi Werneck, 1941
lindfieldi (Hill, 1922)
univirgatus (Neumann, 1913)

Procavicola (Meganarionoides)

africanus Werneck, 1941
angolensis Bedford, 1936

* Known from North America (North of Colombia)

Procavicola (Meganarionoides) (Continued)
colobi (Kellogg, 1910)
congoensis (Ferris, 1930)
jordani Bedford, 1936
muesebecki Emerson and Price, 1969
neumanni baculatus (Ferris, 1930)
neumanni neumanni (Stobbe, 1913)
scutifer Werneck, 1941
tendeiro Emerson, 1964

Procavicola (Procavicola)
affinis Werneck, 1941
brucei Werneck, 1941
eichleri Werneck, 1941
emarginatus (Bedford, 1928)
furca Bedford, 1939
heterohyracis Bedford, 1932
lopesi lopesi Bedford, 1939
lopesi vicinus Werneck, 1941
mokeetsi Bedford, 1939
natalensis Bedford, 1932
parvus Bedford, 1932
pretoriensis Bedford, 1932
shoanus Maltbaek, 1937
sternatus (Bedford, 1928)
subparvus Bedford, 1932
thorntoni Hopkins, 1942
ugandensis Werneck, 1941

Procaviphilus
dubius Werneck, 1941
ferrisi ferrisi Bedford, 1932
ferrisi granuloides Bedford, 1939
ferrisi hindei Werneck, 1946
granulatus (Ferris, 1930)
harrisi Werneck, 1946
robertsi (Bedford, 1928)
sclerotis major Maltbaek, 1937
sclerotis sclerotis Bedford, 1932
serraticus (Hill, 1922)

Stachiella
divaricata (Harrison, 1915)
ermineae Hopkins, 1941 *
jacobi Eichler, 1941

* Known from North America (North of Colombia)

Stachiella (Continued)
kingi (McGregor, 1917) *
larseni Emerson, 1962 *
mustelae (Schrank, 1803)
 (dubius Nitzsch, 1818)
 (pusillus Giebel, 1861)
ovalis (Bedford, 1928)
retusa martis Werneck, 1948 *
retusa retusa (Burmeister, 1818)
retusa salfi Conci, 1940
ugandensis (Bedford, 1936)

Suricatoecus
congoensis Emerson and Price, 1967
cooleyi (Bedford, 1929)
decepiens (Hopkins, 1941)
fahrenheitzi Werneck, 1948
guinlei Werneck, 1948
helogale (Bedford, 1932)
helogaloidis Werneck, 1948
hopkinsi (Bedford, 1936)
laticeps (Werneck, 1942)
mungos (Stobbe, 1913)
paralaticeps Werneck, 1948
quadraticeps (Chapman, 1897) *
vulpis (Denny, 1842)
 (micropus Giebel, 1874)
 (abnormis Ewing, 1930) (?)

Trichodectes
barbarae Neumann, 1913
canis (DeGeer, 1778) *
 (canis O. Fabricius, 1780)
 (latus Nitzsch, 1818)
 (octopunctatus Denny, 1852)
 (riveti Neumann, 1913)
 (floridanus McGregor, 1917)
emersoni Hopkins, 1960
fallax Werneck, 1948 *
ferrisi Werneck, 1944
galictidis Werneck, 1934 *
 (mephitidis Neumann, 1913)
 (paranensis Keler, 1934)
kuntzi Emerson, 1964
malaysianus Werneck, 1948
melis (J. C. Fabricius, 1805)

* Known from North America (North of Colombia)

Trichodectes (Continued)
(crassus Nitzsch, 1818)
octomaculatus Paine, 1912 *
(procyonis Neumann, 1913)
pinguis euarctidos Hopkins, 1954 *
pinguis pinguis Burmeister, 1838
potus Werneck, 1934 *
vosseleri Stobbe, 1913
zorillae Stobbe, 1913

Tricholipeurus
aepycerus Bedford, 1929
albimarginatus (Werneck, 1936) *
annectens (Hopkins, 1943)
antidorcas Bedford, 1931
baculus (Schömmmer, 1913)
bedfordi (Hill, 1922)
clayi (Werneck, 1938)
cornutus balanicus Werneck, 1938
cornutus cornutus (Gervais, 1844)
(antilopes Gurlt, 1857)
cornutus ourebiae (Hopkins, 1943)
dorcelaphi (Werneck, 1936)
elongatus Bedford, 1934
indicus Werneck, 1950
lerouxi Bedford, 1930
lineatus (Bedford, 1920)
lipeuroides (Megnin, 1884) *
(mexicanus Mjöberg, 1910)
(mazama Stobbe, 1913)
(virginianus Peters, 1930)
longiceps (Rudow, 1866)
moschatus Emerson and Price, 1971
parkenhami Werneck, 1947
parallelus (Osborn, 1896) *
(odocoilei McGregor, 1917)
parkeri Hopkins, 1941
spinifer (Hopkins, 1943)
trabeculae reduncae Bedford, 1929
trabeculae trabeculae Bedford, 1929
trabeculae ungangandae Werneck, 1950
victoriae (Hopkins, 1943)

* Known from North American (North of Colombia)

TRIMENOPONIDAE

Chinchillophaga
clayae Emerson, 1964

Cummingsia
inopinata Mendez, 1971
intermedia Werneck, 1937;
maculata Ferris, 1922
peramydis Ferris, 1922
(horridum Harrison, 1922)

Harrisonia
uncinata Ferris, 1922 *

• Hoplomyophilus
nativus Mendez, 1967 *

Philandesia
chinchillae (Werneck, 1935)
mazzai (Werneck, 1933)
townsendi Kellogg and Nakayama, 1914

Trimenopon
hispidum (Burmeister, 1838) *
(echinoderma Cummings, 1913)
(jenningsi Kellogg and Paine, 1910)
(rozeboomi Emerson, 1940)

* Known from North America (North of Colombia)