

3-1-2019

# Host-parasite associations of the *Cratogeomys fumosus* species group and their chewing lice, *Geomydoecus*

Alex Poppinga  
*University of Northern Iowa*

James W. Demastes  
*University of Northern Iowa*

*See next page for additional authors*

Copyright ©2019 Alex Poppinga, James W. Demastes, Theresa A. Spradling, David J. Hafner, and Mark S. Hafner.

Follow this and additional works at: [https://scholarworks.uni.edu/bio\\_facpub](https://scholarworks.uni.edu/bio_facpub)

Part of the [Biology Commons](#)

*Let us know how access to this document benefits you*

---

## Recommended Citation

Poppinga, Alex; Demastes, James W.; Spradling, Theresa A.; Hafner, David J.; and Hafner, Mark S., "Host-parasite associations of the *Cratogeomys fumosus* species group and their chewing lice, *Geomydoecus*" (2019). *Biology Faculty Publications*. 31.  
[https://scholarworks.uni.edu/bio\\_facpub/31](https://scholarworks.uni.edu/bio_facpub/31)

This Article is brought to you for free and open access by the Department of Biology at UNI ScholarWorks. It has been accepted for inclusion in Biology Faculty Publications by an authorized administrator of UNI ScholarWorks. For more information, please contact [scholarworks@uni.edu](mailto:scholarworks@uni.edu).

---

**Authors**

Alex Poppinga, James W. Demastes, Theresa A. Spradling, David J. Hafner, and Mark S. Hafner

# Host-parasite associations of the *Cratogeomys fumosus* species group and their chewing lice, *Geomydoecus*

ALEX POPINGA<sup>1,2</sup>, JAMES W. DEMASTES<sup>1\*</sup>, THERESA A. SPRADLING<sup>1</sup>, DAVID J. HAFNER<sup>3</sup>, AND MARK S. HAFNER<sup>4</sup>

<sup>1</sup> Department of Biology, University of Northern Iowa, Cedar Falls, IA 50614-0421. Email: [alexandrapopinga@gmail.com](mailto:alexandrapopinga@gmail.com) (ANP), [jim.demastes@uni.edu](mailto:jim.demastes@uni.edu) (JWD), [theresa.spradling@uni.edu](mailto:theresa.spradling@uni.edu) (TAS).

<sup>2</sup> Present address: Department of Physics and Department of Computer Science, University of Auckland, Auckland, New Zealand 1010.

<sup>3</sup> Museum of Southwestern Biology, University of New Mexico, Albuquerque, NM 87131, USA. Email: [dhafner@unm.edu](mailto:dhafner@unm.edu) (DJH).

<sup>4</sup> Museum of Natural Science and Department of Biological Sciences, Louisiana State University, Baton Rouge, LA 70803, USA. Email: [namark@lsu.edu](mailto:namark@lsu.edu) (MSH).

\*Correspondence

Chewing lice (Phthiraptera: Trichodectidae) and the pocket gophers (Rodentia: Geomyidae) that they inhabit have shared an intimate historical association that has made them a textbook study for cophylogeny. Herein, we examine the chewing lice found on pocket gophers of the *Cratogeomys fumosus* species group using DNA sequence data from the mitochondrial cytochrome c oxidase subunit I gene to determine relationships among lice for comparison to published host phylogeny. Although matrix approaches indicate a correlation between host and parasite genetic distances, cophylogenetic reconstruction methods fail to detect a pattern of widespread cophylogeny. In conclusion, this study provides an exception to the rule of host-parasite cophylogeny that could be the result of the young age of the relationships considered herein and the complex history of the Trans-Mexican Volcanic Belt.

Los piojos masticadores (Phthiraptera: Trichodectidae) y las tuzas (Rodentia: Geomyidae) que habitan, han compartido una asociación histórica íntima que los ha convertido en un estudio de libro de texto para la cofilogenia. En el presente estudio, examinamos los piojos masticadores que se encuentran en las tuzas del grupo de especies de *Cratogeomys fumosus* utilizando los datos de secuencias de ADN del gen mitocondrial Citocromo C Oxidasa subunidad I para determinar las relaciones entre los piojos en comparación con la filogenia publicada del hospedero. Si bien los enfoques matriciales indican una correlación entre las distancias genéticas del hospedero y el parásito, los métodos de reconstrucción coflogenéticos no detectan un patrón de cofilogenia generalizada. En conclusión, este estudio, proporciona una excepción a la regla de cofilogenia parásito-hospedero que podría ser el resultado de la corta edad de las relaciones consideradas en este documento y la compleja historia del Cinturón Volcánico Trans-Mexicano.

**Key Words:** Cophylogeny; *Cratogeomys*; Geomyidae; mtDNA; Phthiraptera; Trans-Mexican Volcanic Belt.


© 2019 Asociación Mexicana de Mastozoología, [www.mastozoologiamexicana.org](http://www.mastozoologiamexicana.org)

## Introduction

Pocket gophers (Rodentia: Geomyidae) and their chewing lice (Phthiraptera: Trichodectidae) represent a model system for the study of cophylogeny (Hafner *et al.* 2003, and references therein; Light and Hafner 2007). Geomyids are fossorial and asocial, occurring in patchily distributed populations, with genetically distinct groups being largely parapatric (Daly and Patton 1990; Hafner *et al.* 2003). Dozens of non-mammalian organisms inhabit the closed burrow system of pocket gophers and, like the pocket gophers, these organisms are also specialized in this unique niche. In fact, many of these parasites and inquilines are found nowhere else (Hubbel and Goff 1940; Tishechkin and Cline 2008). Among these pocket gopher-dependent organisms are over 100 named species and subspecies of chewing lice (Phthiraptera: Trichodectidae). The majority of these chewing lice seem to be highly host-specific, occurring on a single pocket gopher species or subspecies (Hellenthal and Price 1991). Chewing lice are wingless insects that feed on skin detritus of their hosts (Marshall 1981). The very specializations that make chewing lice well suited for a subterranean existence on a solitary host also greatly reduce their ability to disperse (Marshall 1981). When this poor disper-

sal ability is combined with the solitary nature of geomyids, the probability of colonizing a new host (host switching) is low (Hafner *et al.* 2003). Hence, chewing lice and their pocket gopher hosts are intimately associated. This intimate association continues across evolutionary timescales (Hafner *et al.* 1994; Light and Hafner 2007), making the pocket gopher-lice system a “textbook case” of cophylogeny (*e. g.*, see Page and Holmes 1998; Ridley 2004; Futuyma 2005; Morris *et al.* 2016).

Although the Geomyidae have a distribution that stretches from Canada to northern Colombia (Hafner 2015), the family’s highest diversity (five genera) is found in the Trans-Mexican Volcanic Belt (TMVB). The TMVB consists of hundreds of volcanic structures stretching east to west within the Mexican Transition Zone between the Nearctic-Neotropical realms (Figure 1; Mastretta-Yanes *et al.* 2015). Aside from the striking physiographic relief of this 1,200 km belt, this region is characterized by a long history of volcanism beginning in the Miocene with major structural changes continuing through the Pleistocene and very recent formation of major stratovolcanoes (García-Palomo *et al.* 2000). In addition to this complex geological history, there have been at least three Pleistocene glacial-intergla-



**Figure 1.** Map depicting distribution of the *Cratogeomys fumosus* species group and the five genetic clades delineated by Hafner *et al.* (2004). Symbols indicate localities and *Geomydoecus* louse species sampled for this study.

cial cycles (White 1986; Nixon 1989; Heine 1994a 1994b). These co-occurring events affected community composition and species phylogeography throughout the Pleistocene, and any genetic structure that came about during this period was likely the result of geology, climate, or more likely both (Mastretta-Yanes *et al.* 2015). Given the complex geologic and climatic history, one would expect a complex history of evolution in organisms distributed across this highly plastic landscape, and the genus *Cratogeomys* certainly meets this expectation. Prior to Hafner *et al.* (2004), six species of *Cratogeomys* were recognized from the area encompassed by the TMVB, distributed in a mosaic (Russell 1968). *Cratogeomys merriami* occupied territories from around Mexico City eastward to the east coast. Hafner *et al.* (2004) revised this to include three species comprising the *C. merriami* species group. Russell (1968) referred to *Cratogeomys* in the western half of the TMVB as the *C. gymnurus* species-group, and it was this species-group that most-reflected the complex history of the region with five named species and disjunct, mosaic distributions within *C. tylosinus* and *C. gymnurus*. The complex geologic history, distribution of soils, and plasticity of the morphology in geomyids (Patton and Brylski 1987) undoubtedly made earlier morphological analyses problematic. Hafner *et al.* (2004) revised the *C. gymnurus* species group using to include two species, *C. fumosus* and *C. planiceps* comprising the *C. fumosus* species group (Figure 1). *Cratogeomys planiceps* is found in the vicinity

of Nevado de Toluca Volcano and Valle de Bravo in central Mexico, and *C. fumosus* is broadly distributed across the western half of the TMVB and is composed of four genetically delineated clades that reflect the regions rich history (clades A-D; Figure 1).

The known distribution of the chewing lice hosted by the *C. fumosus* species group is based on the definitive work of Price and Hellenthal (1989). The complex geologic and climatic history of the region along with likely contact with *C. merriami* has resulted in an interesting host-parasite assemblage involving three louse species complexes: the *Geomydoecus mexicanus* and *G. coronadoi* complexes (both also found on the *C. merriami* species group), and the *G. mcgregori* complex (shared with pocket gophers of the genus *Pappogeomys*). Of the five species of chewing lice included in this study, only *G. traubi* and *G. merriami* are also found on other taxa (both occur on *C. merriami*). *Geomydoecus mcgregori* is distributed across the western TMVB and is found on pocket gophers belonging to clades D and C (Price and Hellenthal 1989; Figure 1). *Geomydoecus wernecki* (a member of the *G. mcgregori* species complex) can be found on the remaining gophers of the *C. fumosus* species group (Figure 1). *Geomydoecus polydentatus* is a widely distributed taxon largely sympatric with chewing lice of the *G. mcgregori* species complex. One additional louse taxon, *G. tellus* was described from the northwestern extremes of the distribution of *C. fumosus* (e.g., locality 1, Figure 1; Price and Hellenthal 1988). This louse was morphologically dis-



tinct and has not been assigned membership to any existing species complex.

[Paterson et al. \(2003\)](#) discussed the kinds of historical events that will affect the degree of congruence between phylogenies of associated hosts and parasites. Whereas codivergence yields congruent phylogenies, four types of events can result in varying degrees of phylogenetic incongruence: 1) parasite duplication (speciation of parasite without speciation of host). 2) Parasite inertia (in which the host lineage diverges and the parasite lineage does not). 3) Host switching. 4) Lineage sorting. The asocial nature of pocket gophers and low vagility of chewing lice reduces the likelihood of host switching in this system, whereas lineage sorting is an intrinsic property of any host-parasite interaction. Gene trees that do not match species trees for the parasite lineage, the host lineage, or both can obscure any potential underlying pattern of cophylogeny. Moreover, an ancestral host population may carry multiple genetically divergent lineages of parasites, which can then be retained or lost in a stochastic manner on isolated host populations.

[Demastes et al. \(2012\)](#) demonstrated the effects that differing evolutionary scale can have on the presence of detectable cophylogeny in *Pappogeomys* and their chewing lice. Recently diverged lineages of hosts and parasites are less likely to show cophylogeny than older lineages because recently diverged populations of hosts are more likely to share gene flow (reticulate evolution of the host populations), which increases the likelihood of host switching and detracts from the likelihood of observing parallel phylogenies ([Nieberding and Olivieri 2007](#)). Successful transfer of parasites is more likely in hybridization events between intraspecific populations of hosts because the hosts are more similar (providing similar habitats for the parasites) and because there has been less time for the evolution of host specificity in the parasites. Incomplete lineage sorting also may obscure an underlying pattern of cophylogeny. This is a potential problem for phylogenetic analyses of recently diverged taxa, particularly when effective population size is large ([Maddison and Knowles 2006](#)). Because younger lineages have had less time for lineage sorting of parasites to occur, and because the genes we use to infer relationships in each lineage have had less time to coalesce, we are less likely to observe a pattern of similar evolutionary histories based on genetic analysis of more recently diverged pairs of host and parasite taxa ([Hafner and Page 1995](#); [Rannala and Michalakis 2003](#); [Nieberding and Olivieri 2007](#)).

Like [Demastes et al. \(2012\)](#), other studies of pocket gophers and chewing lice and of Neotropical figs and their pollinators have both shown less cospeciation at finer phylogenetic scales in lineages that appear to cospeciate on a larger scale ([Demastes and Hafner 1993](#); [Demastes et al. 2003](#); [Jackson et al. 2008](#)). These observations, however, are far from exhaustive. As [Huysse et al. \(2005\)](#) suggest, studies of multiple host-parasite systems at multiple taxonomic levels are needed before drawing any conclusions regard-

ing the relationship between taxonomic level and likelihood of cospeciation.

Herein, we examine the chewing lice found on pocket gophers of the *Cratogeomys fumosus* species group using DNA sequence data from the mitochondrial cytochrome c oxidase subunit I (COI) gene. We use these data to compare the resulting genetic distance matrices and phylogenies for these lice with those of their hosts ([Hafner et al. 2004](#)) and test for the presence of a widespread pattern of cophylogeny.

## Materials and Methods

**Specimens examined.** We examined a total of 21 specimens of *Geomydoecus* collected from the same specimens of pocket gophers (*Cratogeomys*), from 19 localities analyzed by [Hafner et al. \(2004\)](#). Specimens were collected under SEMARNAT Permit FAUT-0002 issued to F. A. Cervantes. Following DNA isolation, voucher specimens were preserved following [Cruickshank et al. \(2001\)](#) and identified to species based on the taxonomic characters of [Price and Emerson \(1971\)](#) and [Price and Hellenthal \(1989\)](#).

**Phylogenetic Analysis.** DNA extraction, amplification, and sequencing for chewing lice followed [Light and Hafner \(2007\)](#). Genomic DNA was extracted (DNeasy Tissue Kit, QIAGEN, Valencia, California) from individual chewing lice following the manufacturer's protocol, with a final elution of 30  $\mu$ l for each sample. Extractions were amplified by polymerase chain reaction (PCR) for overlapping regions of the mitochondrial cytochrome c oxidase subunit I (COI) gene, using primers LCO1490 and HCO2198 from [Folmer et al. \(1994\)](#) with an annealing temperature of 46 C for 40 cycles. Two additional primers were designed for this study, HCO758 (AAGAATATATACCTCTGGGTGACC) and HCO740 (GACCAAAAAACCAAAACAAATGC) to be used in conjunction with LCO1490. Success of PCR reactions was determined using agarose gel electrophoresis, and negative PCR controls were used to help detect any possible contamination that may have occurred pre-PCR. Resulting amplified fragments were prepared for sequencing using EXOSAP-it (USB, Cleveland, Ohio), and sequencing was performed at Iowa State University's DNA Facility (Ames, Iowa) using their ABI 3730 DNA Analyzer (Applied Biosystems, Foster City, California). All sequenced regions received at least 2x coverage by sequencing with both PCR primers. Sequences were aligned and edited using Geneious version 11.0 ([Kearse et al. 2012](#)). All sequences were subjected to a BLAST analysis using BLASTN version 2.8 ([Zeng et al. 2000](#)) to confirm target sequence. Sequences were submitted to GenBank (GenBank Accession Nos. KF005296 to KF005316). *Geomydoecus bulleri* (JF342595; from the *G. bulleri* species complex) was used as an outgroup to root all louse trees.

JModelTest version 2.1 ([Guindon and Gascuel 2003](#); [Darriba et al. 2012](#)) was used to select nucleotide substitution models for maximum-likelihood and Bayesian analyses. Based on AIC (Akaike Information Criterion) criteria, the GTR + I + G model was used with the following model parameters: freqA = 0.40, freqC = 0.22, freqG = 0.13, freqT

= 0.25,  $p$ -inv = 0.53, gamma shape parameter = 1.4. Maximum likelihood analysis was conducted using MEGA version 5.0 (Tamura *et al.* 2011). Support for nodes was estimated using 1,000 bootstrap replicates. MrBayes version 2.2 (Ronquist and Huelsenbeck 2003) was used for Bayesian analyses as implemented in Geneious version 11.0 (Kearse *et al.* 2012). Bayesian analysis consisted of paired runs of 4 Markov-Chain Monte Carlo analyses, each using default settings and iterated for  $1.1 \times 10^6$  generations sampled every 200 generations, discarding the initial 110,000 trees (10% of total) sampled. Convergence was determined by examining the MCMC traces for proper mixing. Phylogenetic trees were visualized using FigTree version 1.4 (Rambaut 2012).

**Tests of cophylogeny.** The overall approach taken to test for cophylogeny followed the methods of Demastes *et al.* (2012) by using both matrix-based and topology-based analyses. Matrix-based tests for cophylogeny are topology-free and compare genetic distances for gene regions in the associated host and parasite taxa. Host genetic distances (uncorrected) were calculated from the 1,133 bp from the cytochrome *b* gene sequences from Hafner *et al.* (2004; Genbank AF302155, AF302156, AF302162, AF302163, AF302166-AF302170, AF302174, AF302175, AF302178-AF302183, AY545531-AY545533, AY545535, AY545539-AY545541) and compared to the uncorrected distances calculated for the chewing lice. Both distance matrices were calculated using the program MEGA 5.0 (Tamura *et al.* 2011). The null hypothesis of random association between the host and parasite distance matrices was tested using Mantel tests as implemented in the *ape*, version 5.0, package in R (R Core Team 2014; Paradis and Schliep 2018). Probability of a nonrandom association between the matrices was calculated using the standardized statistic ( $r$ ) with 1,000 permutations. Tests were run on the complete 21 taxon dataset. Further tests were run with matrices that were pruned to only include chewing lice from the *G. mcgregori* species complex and their pocket gopher hosts. A second distance-based method, Parafit (Legendre *et al.* 2002), also was used to compare the distance matrices after converting them into principal component matrices. Tests of random association were conducted with 999 permutations globally and across all pairs of taxa. This analysis was performed using the Parafit option in the *ape*, version 5.0, package in R. Parafit tests were run on the complete 21 taxon dataset, and additional tests were run with matrices that were pruned to only include chewing lice from the *G. mcgregori* species complex and their pocket gopher hosts.

For topology-based cophylogenetic analyses, the host phylogeny was taken from Hafner *et al.* (2004) and pruned to include taxa hosting chewing lice from the *Geomydoecus* species complexes included in this study. The phylogenies of the hosts and parasites were compared to test for a history of widespread cophylogeny. Reconciliation analysis was performed using Jane version 4.0 (Conow *et al.* 2010) with a cost of zero for codivergence events and a cost of one for host switches, duplications, or losses. Significance

was determined by randomization of the parasite tree ( $n = 10,000$ ) and comparison of the resulting null distribution of codivergence events to number of codivergence events estimated from comparison of the data-based host and parasite trees. Analyses were performed on the full 21-taxon phylogeny and on subsets of the chewing louse species complexes.

## Results

Because of the documented presence of at least two different louse species complexes at each of the sampled localities, there was the potential to obtain 38 distinct geographic samples (2 x 19 localities; Table 1). Despite examining multiple lice from each locality (and often multiple hosts individuals), we were only successful in identifying both possible lice from two localities, both within the *C. planiceps* distribution (Table 1). Therefore, at most sites, one of the two possible co-occurring species was absent from the host examined (Table 1).

**Louse Phylogeny.** For the 21 louse individuals included in this study, the 550 bp of COI sequence were generated and analyzed phylogenetically. The aligned dataset held 211 parsimony-informative and 329 invariable sites resulting in all 21 ingroup taxa possessing unique haplotypes. Maximum likelihood and Bayesian analyses produced phylogenetic trees with nearly identical topologies. Minor differences between maximum-likelihood and Bayesian trees for the lice disappeared when poorly supported nodes (less than 70 % bootstrap support or less than 0.8 posterior probability) were collapsed, producing a phylogeny that is largely concordant with current louse taxonomy (Figure 2). Genetic data revealed the presence of two genetically distinct lineages within *G. mcgregori*. Some of the louse clades recovered (Figure 2) are consistent with both geographic proximity and host taxonomy (Figure 1; *G. mcgregori*, localities 1, 2, and 4 and *G. wernecki*, localities 8, 9, and 11). However, one louse clade (*G. mcgregori*, localities 3, 5, and 7) includes geographic neighbors derived from hosts belonging to different genetic groups. Another louse clade (*G. wernecki*, localities 10 and 19) includes specimens that are not in geographic proximity and that occur on hosts belonging to different genetic groups. Likewise, other closely related lice (for example, *G. polydentatus*, localities 12 and 21 and *G. traubi* localities 16 and 20) are found on geographically distant hosts belonging to different genetic groups.

**Tests of Cophylogeny.** Matrix comparison for genetic distances of chewing lice and pocket gophers indicated a significant correlation for both the complete dataset (Mantel test,  $z = 3.19$ ,  $P = 0.007$ ) and for matrices trimmed to include only the clades involving the *G. mcgregori* complex ( $z = 0.68$ ,  $P = 0.005$ ). Similar results were returned using the Parafit algorithm. Global values for the complete dataset ( $F = 0.006$ ,  $P = 0.008$ , 10 of 21 links significant) and the trimmed version ( $F = 0.007$ ,  $P = 0.004$ , 7 of 12 links significant) also indicated a significant positive correlation for the host and parasite distance matrices.

**Table 1.** Specimens examined. Map column refers to Figure 1. “*Geomydoecus* complex” indicates which chewing louse species could be expected from each species complex on the *Cratogeomys* host listed compared with locality data from Price and Hellenthal (1989). Asterisks indicate the louse species actually identified and included in this study. Note that several additional lice from the same and from different hosts were examined from each locality but, to date, in no case have two species been found on the same host. LSUMZ = Louisiana State University, Museum of Natural Science.


Map	Host	Collection #	Geographic Locality	Geomydoecus complex		
				<i>mcgregori</i>	<i>mexicanus</i>	<i>coronodoi</i>
1	<i>C. fumosus</i>	LSUMZ 36081	Jalisco: 16 km E Ameca <sup>1</sup>	<i>mcgregori</i> *		<i>polydentatus</i>
2	<i>C. fumosus</i>	LSUMZ 36077	Jalisco: 5 km SW Mazamitla	<i>mcgregori</i> *		<i>polydentatus</i>
3	<i>C. fumosus</i>	LSUMZ 36164	Michoacán: 5 km N Tingüindín	<i>mcgregori</i> *		<i>polydentatus</i>
4	<i>C. fumosus</i>	CNMA 39673	Michoacán: 2 km N, 5 km W Apo, 1,720 m	<i>mcgregori</i> *		<i>polydentatus</i>
5	<i>C. fumosus</i>	LSUMZ 36133	Michoacán: 2 km N Uruapan	<i>mcgregori</i> *		<i>polydentatus</i>
6	<i>C. fumosus</i>	LSUMZ 36292	Michoacán: 2 km NW Patamban	<i>mcgregori</i>		<i>polydentatus</i> *
7	<i>C. fumosus</i>	LSUMZ 36124	Michoacán: Tangancicuaro	<i>mcgregori</i> *		<i>polydentatus</i>
8	<i>C. fumosus</i>	LSUMZ 34425	Michoacán: 6.5 km S Pátzcuaro	<i>wernecki</i> *		<i>polydentatus</i>
9	<i>C. fumosus</i>	LSUMZ 36136	Michoacán: San Gregorio, 8 km E Opopeo	<i>wernecki</i> *		<i>polydentatus</i>
10	<i>C. fumosus</i>	LSUMZ 36130	Michoacán: 5 km S, 20 km E Morelia	<i>wernecki</i> *		<i>polydentatus</i>
11	<i>C. fumosus</i>	LSUMZ 36129	Michoacán: 1 km S Tacámbaro	<i>wernecki</i> *		<i>polydentatus</i>
12	<i>C. fumosus</i>	LSUMZ 36092	Guanajuato: 1 km E Celaya	<i>wernecki</i>		<i>polydentatus</i> *
13	<i>C. fumosus</i>	LSUMZ 36091	Querétaro: La Cañada, 9 km (by road) SW Pinal de Amoles	<i>wernecki</i>		<i>polydentatus</i> *
14	<i>C. planiceps</i>	LSUMZ 36075	México: 25 km N Valle de Bravo		<i>traubi</i> *	<i>merriami</i>
15	<i>C. planiceps</i>	LSUMZ 36303	México: 25 km N Valle de Bravo		<i>traubi</i>	<i>merriami</i> *
16	<i>C. planiceps</i>	LSUMZ 36123	México: 3 km S, 20 km E Valle de Bravo		<i>traubi</i> *	<i>merriami</i>
17	<i>C. planiceps</i>	LSUMZ 36291	México: 3 km S, 20 km E Valle de Bravo		<i>traubi</i>	<i>merriami</i> *
18	<i>C. planiceps</i>	LSUMZ 34901	México: 10 km S, 16 km W Toluca, 3,000 m	<i>wernecki</i>	<i>traubi</i> *	
19	<i>C. fumosus</i>	LSUMZ 36163	México: 26 km W Toluca	<i>wernecki</i> *	<i>traubi</i>	
20	<i>C. fumosus</i>	LSUMZ 34902	México: 1 km N La Isla, 2,612 m	<i>wernecki</i>	<i>traubi</i> *	
21	<i>C. fumosus</i>	LSUMZ 36066	México: 1 km S Tepexpan	<i>wernecki</i>		<i>polydentatus</i> *

<sup>1</sup> The morphologically unique *Geomydoecus tellus* has also been described from this locality and co-occurring on the same host individual as *G. mcgregori* (Price and Hellenthal, 1988). It has not been assigned to a species complex and was not detected in our sampling.

Tanglegram comparison superficially indicated little similarity between host and parasite phylogenies (Figure 3). Cophylogenetic reconstruction analysis using Jane (Conow et al. 2010) indicated the absence of a widespread pattern of cophylogeny for the chewing lice and pocket gophers. In that analysis, twenty-eight percent of the cophylogenetic reconstructions using the Hafner et al. (2004) gopher topology and a randomly generated topology for the chewing lice had costs that were equal to or less than the cost of the best (lowest cost) reconstruction for the observed data (cost of observed topologies = 16;  $P = 0.3$ ). Analyses were repeated for each chewing louse species complex (Figure 4). These analyses for the *G. mcgregori*, *G. coronadoi*, and *G. mexicanus* complexes also indicated the lack of a widespread nonrandom pattern of cophylogeny ( $P = 0.3$ ,  $P = 0.5$ , and  $P = 0.7$  respectively). However, some of these comparisons were hindered by small sample sizes and, short of perfect congruence, were not likely to indicate statistical significance.

### Discussion

**Louse Phylogeny.** Past genetic examinations of morphologically-based phylogenies of chewing lice of pocket gophers have revealed a remarkable level of accuracy in the earlier taxonomic studies, even at the subspecies level (e. g., Hafner and Nadler 1988; Demastes and Hafner 1993; Light



**Figure 2.** Maximum likelihood (ML) tree for 21 *Geomydoecus* lice studied (*G. mcgregori* in inset). Locality numbers refer to Figure 1 and Table 1. Numbers on branches indicate ML bootstrap support/Bayesian posterior probability.




and Hafner 2007). Our molecular data indicated genetically cohesive groups of individuals consistent with the morphologically-based species-level taxonomy of Price and Hellenenthal (1989), with the exception that they reveal for the first time the existence of two divergent, well-supported and potentially paraphyletic clades of lice within *G. mcgregori*. Although these two *G. mcgregori* clades (localities 1, 2, 4 and localities 3, 5, 7) are not concordant with their associated host phylogenies, they do occur as geographically cohesive units (Figure 1).

**Tests of Cophylogeny.** As reviewed by de Vienne et al. (2013), there are distinct advantages and disadvantages to both matrix-based and reconciliation-based methods for the testing of cophylogeny. One drawback for matrix-based methods such as Mantel tests and ParaFit is their inability to account for phylogenetic non-independence (Felsenstein 1985). This in itself could account for a possible false-positive signal of cospeciation from matrix-based approaches. While matrix-based approaches indicated significant congruence between louse and host genetic distances in this study, the more rigorous reconciliation analyses suggested no significant pattern of cophylogeny. Relatively few nodes are shared between host and parasite trees, indicating a large degree of discordance between host and parasite phylogenies, with several closely related louse specimens occurring on rather distantly related hosts (Figure 3).

Past studies have established that pocket gophers and chewing lice frequently show a pattern of cospeciation, an outcome likely favored by the life histories of both lineages


(e. g., Hafner et al. 2003; Light and Hafner 2007; Demastes et al. 2012). The pattern of non-cospeciation observed here may be the result of several factors. First, results in this study come from a portion of a single mitochondrial gene. While the pivotal nodes of the parasite tree are well supported in our analyses, it is possible that a more robust analysis with additional genetic loci from the nucleus could reveal a different pattern of relationships that reflect an underlying pattern of cospeciation if incomplete lineage sorting in the mitochondrial data has incorrectly reflected the true phylogenetic history of the group. Secondly, several potential pairs of co-occurring chewing lice were not sampled, either because they are rare or because they have been extirpated from the sampled localities. Loss of these populations could obscure a pattern of cophylogeny if extirpation was followed by colonization by a new form. The *G. wernecki* at locality 3 (Tingüindín, Michoacán) may be the result of such an extirpation/colonization event given that the phylogenetic position of this louse (Figure 3) is more consistent with geography than with host relationships (Figure 1). It is also possible that the recent age of this particular system may play a role in the lack of cospeciation observed here by facilitating the likelihood of successful host switching and/or through the influence of lineage-sorting effects (Hafner and Page 1995; Rannala and Michalakis 2003; Nieberding and Olivieri 2007; Demastes et al. 2012).

Finally, as discussed previously (and in detail in Demastes et al. 2002; pp. 150-152) the Quaternary history of the western TMVB is quite complex with major geologic



**Figure 3.** Tanglegram illustrating host parasite associations between pocket gophers (*Cratogeomys*) and their chewing lice (*Geomydoecus*). The pocket gopher phylogeny is taken from Hafner et al. (2004) and pruned to include only pertinent specimens.





**Figure 4.** Subtree tanglegrams for louse species complexes and their associated pocket gopher hosts.

events occurring after estimated times of speciation for the hosts included in this study, *C. fumosus* and *C. planiceps*. The distributions for the five differentiated clades of pocket gophers included here (Hafner et al. 2004) likely experienced major changes during periods of increased volcanism and climatic shifts in the Pleistocene and Holocene. This complex history of shifting distributions caused by natural events would have presented many opportunities for contact between host individuals of different genetic groups and subsequent host switching of parasites (Pater-son et al. 2003). The complex, dynamic history of the western TMVB has been the driving force behind the generation of the unique diversity of the region (González-Fernández et al. 2018; Mastretta-Yanes et al. 2015; Navarro-Sigüenza et al. 2007). As more phylogeographic studies are conducted on co-distributed taxa of the TMVB, a comparative approach may provide greater insights into the effects these events had on the biota of the region. For example, an alligator lizard (genus *Barisia*) exhibits a pattern of geographic differentiation that is very similar to that of *Cratogeomys* across the TMVB (Bryson and Riddle 2012). It is possible that frequent range changes resulting from climate change in this geologically dynamic region have generated an assemblage of

pocket gophers and chewing lice with frequent opportunities for host-switching. This, coupled with a great potential for isolation and local extinction of chewing lice in the geographically intricate TMVB, could have caused obscured the normal pattern of cophylogeny typically observed for geomyids and their trichodectid lice.

### Acknowledgements

We thank our collaborator in Mexico, F. Cervantes, and his students R. Castro, R. Rojas, J. Castello, and M. Aquino Camacho for valuable assistance in the field and for generous hospitality. J. Light also provided valuable assistance in the field. This research was supported by National Science Foundation grant 0075381 (to M. S. Hafner).

### Literature cited

BRYSON JR, R. W., AND B. R. RIDDLE. 2012. Tracing the origins of widespread highland species: a case of Neogene diversification across the Mexican sierras in an endemic lizard. *Biological Journal of the Linnean Society* 105: 382-394.

CONOW, C., D. FIELDER, Y. OVADIA, AND R. LIBESKIND-HADAS. 2010. Jane: a new tool for the cophylogeny reconstruction problem. *Algorithms for Molecular Biology* 5:16.

- CRUICKSHANK, R. H., K. P. JOHNSON, V. S. SMITH, R. J. ADAMS, D. H. CLAYTON, AND R. D. M. PAGE. 2001. Phylogenetic analysis of partial sequences of elongation factor 1 $\alpha$  identifies major groups of lice (Insecta: Phthiraptera). *Molecular Phylogenetics and Evolution* 19:202–215.
- DARRIBA D., G. L. TABOADA, R. DOALLO, AND D. POSADA. 2012. jModelTest 2: more models, new heuristics and parallel computing. *Nature Methods* 9:772.
- DALY, J. C., AND J. L. PATTON. 1990. Dispersal, gene flow, and allelic diversity between local populations of *Thomomys bottae* pocket gophers in the coastal regions of California. *Evolution* 44:1283–1294.
- DEMASTES, J. W., AND M. S. HAFNER. 1993. Cospeciation of pocket gophers (*Geomys*) and their chewing lice (*Geomydoecus*). *Journal of Mammalogy* 74:521–530.
- DEMASTES, J. W., T. A. SPRADLING, M. S. HAFNER, D. J. HAFNER, AND D. L. REED. 2002. Systematics and phylogeography of pocket gophers in the genera *Cratogeomys* and *Pappogeomys*. *Molecular Phylogenetics and Evolution* 2:144–154.
- DEMASTES, J. W., T. A. SPRADLING, AND M. S. HAFNER. 2003. The effects of spatial and temporal scale on analyses of cophylogeny. Pp. 221–239 in *Tangled trees: Phylogeny, cospeciation, and coevolution* (Page, R. D. M. ed.). The University of Chicago Press, Chicago, U. S. A.
- DEMASTES, J. W., T. A. SPRADLING, M. S. HAFNER, G. R. SPIES, D. J. HAFNER, AND J. E. LIGHT. 2012. Cophylogeny on a fine scale: *Geomydoecus* chewing lice and their pocket gopher hosts, *Pappogeomys bulleri*. *Journal of Parasitology* 98:262–270.
- DE VIENNE, D. M., G. REFRÉGIER, M. LÓPEZ-VILLAVICENCIO, A. TELLIER, M. E. HOOD, AND T. GIRAUD. 2013. Cospeciation vs host shift speciation: methods for testing, evidence from natural associations and relation to coevolution. *New Phytologist* 198:347–385.
- FELSENSTEIN, J. 1985. Phylogenies and the comparative method. *The American Naturalist* 125:1–15.
- FOLMER, O., M. BLACK, W. HOEH, R. LUTZ, AND R. VRIJENHOEK. 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Molecular Marine Biology and Biotechnology* 3:294–299.
- FUTUYMA, D. J. 2005. *Evolution*. Sinauer Associates, Sunderland, Massachusetts, U. S. A.
- GARCÍA-PALOMO, A., J. L. MACÍAS, AND V. H. GARDUÑO. 2000. Miocene to recent structural evolution of the Nevado de Toluca volcano region, Central Mexico. *Tectonophysics* 318:281–302.
- GONZÁLEZ-FERNÁNDEZ, A., J. MANJARREZ, U. GARCÍA-VÁZQUEZ, M. D'ADDARIO, AND A. SUNNY. 2018. Present and future ecological niche modeling of garter snake species from the Trans-Mexican Volcanic Belt. *PeerJ* 6:e4618.
- GUINDON, S., AND O. GASCUEL. 2003. A simple, fast and accurate method to estimate large phylogenies by maximum-likelihood. *Systematic Biology* 52:696–704.
- HAFNER, M. S. 2015. Family Geomyidae, Bonaparte, 1845. Pp 49–51 in *Mammals of South America, Volume 2*, (J. L. Patton, U. F. Pardiñas, and G. D'Elia, eds.). University of Chicago Press, Chicago, Illinois, U. S. A.
- HAFNER, M. S., AND S. A. NADLER. 1988. Phylogenetic trees support the coevolution of parasites and their hosts. *Nature* 332:258.
- HAFNER, M. S., P. D. SUDMAN, F. X. VILLABLANCA, T. A. SPRADLING, J. W. DEMASTES, AND S. A. NADLER. 1994. Disparate rates of molecular evolution in cospeciating hosts and parasites. *Science* 265:1087–1090.
- HAFNER, M. S., AND R. D. M. PAGE. 1995. Molecular phylogenies and host–parasite cospeciation: Gophers and lice as a model system. *Philosophical Transactions of the Royal Society, B* 349:77–83.
- HAFNER, M. S., J. W. DEMASTES, T. A. SPRADLING, AND D. L. REED. 2003. Cophylogeny in pocket gophers and chewing lice. Pp. 195–220 in *Tangled trees: Phylogeny, cospeciation, and coevolution* (Page, R. D. M. ed.). The University of Chicago Press, Chicago, U. S. A.
- HAFNER, M. S., T. A. SPRADLING, J. E. LIGHT, D. J. HAFNER, AND J. R. DEMBOSKI. 2004. Systematic revision of pocket gophers of the *Cratogeomys gymnurus* species group. *Journal of Mammalogy* 85:1170–1183.
- HEINE, K. 1994a. The late-glacial moraine sequences in Mexico: Is there evidence for the younger Dryas event? *Paleogeography, Paleoclimatology, Paleoecology* 112:113–123.
- HEINE, K. 1994b. Past and present geocryogenic processes in Mexico. *Permafrost Periglacial Proceedings* 5:1–12.
- HELLENTHAL, R. A., AND R. D. PRICE. 1991. Biosystematics of the chewing lice of pocket gophers. *Annual Review of Entomology* 36:185–203.
- HUBBEL, T. H., AND C. C. GOFF. 1940. Florida pocket gopher burrows and their arthropod inhabitants. *Proceedings of the Florida Academy of Science* 4:127–166.
- HUYSE, T., R. POULIN, AND A. THÉRON. 2005. Speciation in parasites: A population genetics approach. *Trends in Parasitology* 21:467–475.
- JACKSON, A. P., C. A. MACHADO, N. ROBBINS, AND E. A. HERRE. 2008. Multi-locus phylogenetic analysis of neotropical figs does not support co-speciation with the pollinators: The importance of systematic scale in fig/wasp cophylogenetic studies. *Symbiosis* 45:57–72.
- KEARSE, M., R. MOIR, A. WILSON, S. STONES-HAVAS, M. CHEUNG, S. STURROCK, S. BUXTON, A. COOPER, S. MARKOWITZ, C. DURAN, T. THIERER, B. ASHTON, P. MENTJES, AND A. DRUMMOND. 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28:1647–1649.
- LEGENDRE, P., Y. DESDEVISES, AND E. BAZIN. 2002. A statistical test for host–parasite coevolution. *Systematic Biology* 51:217–234.
- LIGHT, J. E., AND M. S. HAFNER. 2007. Cophylogeny and disparate rates of evolution in sympatric lineages of chewing lice on pocket gophers. *Molecular Phylogenetics and Evolution* 45:997–1013.
- MADDISON, W. P., AND L. L. KNOWLES. 2006. Inferring phylogeny despite incomplete lineage sorting. *Systematic Biology* 55:21–30.
- MARSHALL, A. G. 1981. *The ecology of ectoparasitic insects*. Academic Press. London, U.K.
- MASTRETTA-YANES, A., A. MORENO-LETELIER, D. PINERO, T. H. JORGENSEN, AND B. C. EMERSON. 2015. Biodiversity in the Mexican highlands and the interaction of geology, geography and climate within the Trans-Mexican Volcanic Belt. *Journal of Biogeography* 42:1586–1600.
- MORRIS, J., D. HARTL, A. KNOLL, R. LUE, M. MICHAEL, A. BERRY, A. BIEWENER, B. FARRELL, AND N. M. HOLBROOK. 2016. *Biology: How life works*. W. H. Freeman Co. New York, U. S. A.

- NAVARRO-SIGÜENZA, A. G., A. LIRA-NORIEGA, A. T. PETERSON, A. OLIVERAS DE ITA, AND A. GORDILLO-MARTÍNEZ. 2007. Diversidad, endemismo y conservación de las aves. *Biodiversidad de la Faja Volcánica Transmexicana* 461–483.
- NIEBERDING, C. M., AND I. OLIVIERI. 2007. Parasites: proxies for host genealogy and ecology? *Trends in Ecology and Evolution* 22:156–165.
- NIXON, T. G. 1989. The geology of Iztaccihuatl Volcano and adjacent areas of the Sierra Nevada and Valley of Mexico. *Geological Society of America, Special Paper* 219:1–58.
- PAGE, R. D. M., AND E. C. HOLMES. 1998. *Molecular evolution: a phylogenetic approach*. Blackwell Science Ltd., Oxford, U. K.
- PARADIS, E., AND K. SCHLIEP. 2018. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics* 1:3.
- PATERSON, A. M., R. L. PALMA, AND R. D. GRAY. 2003. Drowning on arrival, missing the boat, and x-events: How likely are sorting events? Pp. 287–309 in *Tangled trees: Phylogeny, cospeciation, and coevolution* (Page, R. D. M. ed.). The University of Chicago Press, Chicago, Illinois, U. S. A.
- PATTON, J. L., AND P. V. BRYLSKI. 1987. Pocket gophers in alfalfa fields: causes and consequences of habitat-related body size variation. *American Naturalist* 130:493–506.
- PRICE, R. D., AND K. C. EMERSON. 1971. A revision of the genus *Geomydoecus* (Mallophaga: Trichodectidae) of the New World pocket gophers (Rodentia: Geomyidae). *Journal of Medical Entomology* 8:228–257.
- PRICE, R. D., AND R. A. HELLENTHAL. 1988. A new species of *Geomydoecus* (Mallophaga: Trichodectidae) from *Pappogeomys* (Rodentia: Geomyidae) pocket gophers in Jalisco, Mexico. *Journal of Entomological Science* 23:212–215.
- PRICE, R. D., AND R. A. HELLENTHAL. 1989. *Geomydoecus* (Mallophaga: Trichodectidae) from *Pappogeomys* and *Zygogeomys* pocket gophers (Rodentia: Geomyidae) in central Mexico. *Journal of Medical Entomology* 26:385–401.
- RAMBAUT, A. 2012. FigTree v. 1.4. <http://tree.bio.ed.ac.uk/software/figtree/>
- RANNALA, B., AND Y. MICHALAKIS. 2003. Population genetics and cospeciation: From process to pattern. Pp. 120–143 in *Tangled trees: Phylogeny, cospeciation, and coevolution* (Page, R. D. M. ed.). The University of Chicago Press, Chicago, U. S. A.
- R CORE TEAM. 2014. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org/>.
- RIDLEY, M. 2004. *Evolution*, Third Ed. Blackwell Publishing, Malden, Massachusetts.
- RONQUIST, F., AND J. P. HUELSENBECK. 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19:1572–1574.
- RUSSELL, R. J. 1968. Revision of pocket gophers of the genus *Pappogeomys*. University of Kansas Publications, Museum of Natural History 16:581–776.
- TAMURA, K., D. PETERSON, N. PETERSON, G. STECHER, M. NEI, AND S. KUMAR. 2011. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular Biology and Evolution* 28:2731–2739.
- TISHECHKIN, A. K., AND A. R. CLINE. 2008. The beetle (Coleoptera) fauna of pocket gopher burrows in Louisiana. *Proceedings of the Entomological Society of Washington* 110:331–339.
- WHITE, S. E. 1986. Quaternary glacial stratigraphy and chronology of Mexico. *Quaternary Science Review* 5:201–205.
- ZHANG, Z., S. SCHWARTZ, L. WAGNER, AND W. MILLER. 2000. A greedy algorithm for aligning DNA sequences. *Journal of Computational Biology* 7:203–214.

Associated editor: Jesús Fernández

Submitted: December 4, 2018; Reviewed: January 22, 2019;

Accepted: February 19, 2019; Published on line: March 1, 2019.

