

## Review

# Lice Infestation in Poultry

MUHAMMAD NISAR KHAN, M. NADEEM, ZAFAR IQBAL, MUHAMMAD SOHAIL SAJID AND RAO ZAHID ABBAS  
*Department of Veterinary Parasitology, University of Agriculture, Faisalabad-38040, Pakistan*

### ABSTRACT

This paper reviews the studies on the prevalence, chemotherapeutic and biological control of lice. It was found that *Menacanthus stramineus*, *Menopon gallinae*, *Goniodes gigas*, *Goniocotes gallinae*, *Lipeurus lawrensis tropicalis*, *Lipeurus caponis* and *Cuclotogaster heterographus* were the most commonly infesting lice of poultry. A variety of chemical are used for the control of lice infestation in poultry. These include Carbamates, Pyrethroids, Cypermethrin, Trichlorphon, Dichlorvos, DDT powder, Coumaphos, Lindane and BHC, Phenol, Cresols, Permethrin, Pestoban, Malathion, Pflspray, Carbaryl, Methyl bromfenvinphos and Herbal products. The most promising results however were reported with Pyrethroids i.e. Trichlophon and Pestoban. Biological control although not widely accepted for application, yet some bacterial and fungal species were found to be effective. Based on the review conclusion have been drawn for further research in to the lice infestation of poultry.

**Key Words:** Lice; Poultry

### INTRODUCTION

Poultry is an economic and effective source of animal protein within the shortest possible time, playing a vital role in narrowing down the animal protein supply gap. Poultry can easily be infected with several types of bacterial, viral, fungal and parasitic pathogens (Soulsby, 1982). Among various parasitic diseases, ectoparasites infestation are of great importance. Ectoparasites of poultry live on the skin or penetrate within the skin or even in to the air sacs and some live under the feathers. These ectoparasites consume dead cells of skin and tissue fluids, while other suck blood (Phillip, 1963; Urquhart, 1987). Lice is an important ectoparasite of poultry which cause ill health in poultry and cause heavy morbidity by sucking blood and causing irritation to the birds, which adversely affects the economical production of poultry (Edgar & King, 1950). The birds reduce haemoglobin and erythrocyte values and hyperchromic anaemia (Jungmann, 1970). Lice infestation causes weight loss at the rate of about 711 gms per birds and decrease the egg yield at the rate of about 66 egg per bird in a year (El-Kifl *et al.*, 1973) and lameness is associated with heavy lice infestation (Okaeme, 1989). The purpose of this paper is to review the studies on lice infestation in poultry and associated risks.

**Prevalence.** There are a number of a species of lice infesting poultry. These include *Menacanthus stramineus* (Hohorst, 1939; Erlich, 1942; Giles & Raun, 1959; Stockdale, 1960; Knapp, 1962; Hoffman & Hogan, 1967; Hoffman & Gingrich, 1968; Brown, 1970; Iqbal *et al.*, 1971; Mathysse, 1972; Mathur, 1973; Martin, 1973; Kumar & Sahai, 1974; Puchkova, 1977; Kachekova & Frolov, 1978; Shafique, 1979; Horning *et al.*, 1980; Reddy *et al.*, 1980; Grzywinski *et al.*, 1980; Zlotorzycska, 1980; Devi & Mishra, 1981; Mehl, 1981; Urban & Zlotorzycska, 1981;

Cariaso, 1982; Modrzejewska, 1982; Zlotorzycska *et al.*, 1982; Panda & Akluwali, 1983; Fedorenko *et al.*, 1983; Hofstad, 1984; Frolov *et al.*, 1984; Reddy *et al.*, 1985; Ratanasethakul *et al.*, 1985; Morrow, 1986; Ugochukwu & Omije, 1986; De-Vaney, 1986; Ahmed, 1986; Bilinski & Jankowska, 1987; Chandra *et al.*, 1988; Okaeme, 1989; Salisch, 1989; Pavlovic *et al.*, 1989; Werner *et al.*, 1989; Noh, Y. T *et al.*, 1989; Ahmad, 1991; George *et al.*, 1992; Chhabra & Donoru, 1994; Shahjehan & Iqbal, 1995; Zhang-Deling *et al.*, 1996; Amin-Babjee *et al.*, 1997; Axtell, 1999; Rafiq, 2000), *Menacanthus cornutus* (Hohorst, 1939; Hohorst, 1942; Erlich, 1942; Mathysse, 1972; El-Kifl *et al.*, 1973; Martin-Mateo *et al.*, 1980; Fabiyi, 1980; Hofstad, 1984; Reddy *et al.*, 1985; Morrow, 1986; Abdul-Karim *et al.*, 1988), *Menacanthus pallidulus* (Hohorst, 1939; Mathysse, 1972; Martin, 1973; Manuel & Anceno, 1981; Hofstad, 1984; Sinha *et al.*, 1989; Noh, Y.T *et al.*, 1989) *Menacanthus cornutum* (Hohorst, 1939; Mathysse, 1972; Martin-Mateo *et al.*, 1980; Fedorenko *et al.*, 1983) *Menacanthus numidae* (Mathysse, 1972; Martin, 1973; Martin-Mateo *et al.*, 1980; Fedorenko *et al.*, 1983), *Cuclotogaster heterographus* (Mathysse, 1972; Kumar & Sahi, 1974; Martin-Marteo *et al.*, 1980; Hofstad, 1984; Reddy *et al.*, 1985; Ahmed, 1986; Morrow, 1986; Shahjehan & Iqbal, 1995; Amin-Babjee *et al.*, 1997), *Cuclotogaster occidentalis* (Fabiyyi, 1980; Matin-Mateo *et al.*, 1980; Mehl, 1981; Modrzejewska, 1982; Fedorenko *et al.*, 1983), *Columbicola colombae* (Hathaway, 1943; Hashmi, 1971; El-Kifl *et al.*, 1973; Mathur, 1973; Singh & Chhabra, 1973; Tigin, 1973; Agarwal & Saxena, 1981; Devi & Mishra, 1981; Mehl, 1981; Dikaev, 1988; Kaminjolo *et al.*, 1988; Noh, Y.T *et al.*, 1989), *Lipeurus lawrensis tropicalis* (Mathysse, 1972; Mathur, 1973; Kumar & Sahai, 1974; Marin-Mateo *et al.*, 1980; Fabiyi, 1980; Zlotorzycska, 1982; Agarwal & Saxena, 1981; Buriro, 1982; Panda &

Akluwali, 1983; Hofstad, 1984; Reddy *et al.*, 1985; Okaeme, 1989; Noh, Y.T *et al.*, 1989), *Lipeurus caponis* (Hohorst, 1939; Erlich, 1942; Hoffman & Hogan, 1967; Hoffman & Gingrich, 1968; Iqbal *et al.*, 1971; Mathysse, 1972; Kumar & Sahai, 1974; Shafique, 1979; Fabiyi, 1980; Devi & Mishra, 1981; Manuel & Aceno, 1981; Buriro, 1982; Hofstad, 1984; Ahmed, 1986; Ugochukwu & Omije, 1986; Okaeme, 1989; Ahmad, 1991; George *et al.*, 1992; Ahmed-Babjee *et al.*, 1997; Islam *et al.*, 1999; Rafiq, 2000), *Lipeurus tropicalis* (Mathur, 1973; Mehl, 1981; Modrzejewska, 1982; Fedorenko *et al.*, 1983; Panda & Akluwali, 1983; Sinha *et al.*, 1989), *Lipeurus heterographus* (Hoyle, 1938; Hohorst, 1939; Martin-Mateo *et al.*, 1980; Mehl, 1981; Modrzejewska, 1982), *Menopon gallinae* (Hoyle, 1938; Erlich, 1942; Hoffman & Hogan, 1967; Hoffman & Gingrich, 1968; Jungmann *et al.*, 1970; Mathysse, 1972; El-Kifl *et al.*, 1973; Martin, 1973; Mathur, 1973; Kumar & Sahai, 1974; Shafique, 1979; Grzywinski *et al.*, 1980; Reddy *et al.*, 1980; Devi & Mishra, 1981; Manuel & Anceno, 1981; Urban & Zlotorzycza, 1981; Buriro, 1982; Cariaso, 1982; Zlotorzycza, 1982; Frolov *et al.*, 1984; Bilqees & Khan, 1985; Ratanasethakul *et al.*, 1985; Ahmad, 1986; Okaeme, 1989; Sinha *et al.*, 1989; Ahmad, 1991; George *et al.*, 1992; Chhabra & Donora, 1994; Sexena *et al.*, 1995; Shahjehan & Iqbal, 1995; Amin-Babjee *et al.*, 1997; Larramendy-R *et al.*, 2000; Rafiq, 2000), *Goniocotes gigas* (Erlich, 1942; Devi & Mishra, 1981; Mahl, 1981; Buriro, 1982; Fedorenko *et al.*, 1983; Frolov *et al.*, 1984; Bilqees & Khan, 1985; Sinha *et al.*, 1989; Shahjehan & Iqbal, 1995) *Goniocotes hologaster* (Hohorst, 1939; Erlich, 1942; Jungmann *et al.*, 1970; Tigin, 1973; Devi & Mishra, 1981; Frolov *et al.*, 1984; Sinha *et al.*, 1989), *Goniocotes gallinae* (Mathysse, 1972; El-Kifl *et al.*, 1973; Mathur, 1973; Ruprah, 1979; Fabiyi, 1980; Grzywinski *et al.*, 1980; Manuel & Anceno, 1981; Devi & Mishra, 1981; Urban & Zlotorzycza, 1981; Zlotorzycza *et al.*, 1982; Hofstad, 1984; Ahmed, 1986; Okaeme, 1989; Chhabra & Donora, 1994; Amin-Babjee *et al.*, 1997), *Goniocotes hidentatus* (Manuel & Anceno, 1981; Buriro, 1982; Fedorenko *et al.*, 1983), *Goniocotes bidentatus* (Tigin, 1973; Singh & Chhabra, 1973; Fedorenko *et al.*, 1983), *Goniodes gigas* (Mathysse, 1972; Mathur, 1973; Kumar & Sahai, 1974; Fabiyi, 1980; Hofstad, 1984; Ratanasethakul *et al.*, 1985; Ugochukwu & Omije, 1986; Okaeme, 1989; Ahmad, 1991; George *et al.*, 1992; Chhabra & Donora, 1994; Amin-Babjee *et al.*, 1997; Rafiq, 2000) *Goniodes dissimilis* (Hohorst, 1939; Mathysse, 1972; Kumar & Sahai, 1974; Fabiyi, 1980; Manuel & Anceno, 1981; Shusmanou, 1982; Fedorenko *et al.*, 1983; Hofstad, 1984; Ratanasethakul *et al.*, 1985; Noh, Y.T *et al.*, 1989; Shahjehan & Iqbal, 1995; Amin-Babjee *et al.*, 1997), *Goniodes pavonis* (Martin-Mateo *et al.*, 1980; Rodriguez *et al.*, 1981; Fedorenko *et al.*, 1983; Okaeme, 1989)

Incidence rates remained distinctly higher on older birds, the intensity of infection remained distinctly higher during the summer months (Saxena *et al.*, 1995). The major out brakes of lice infestation occurred in the hot dry

period of February-May (Fabiyi, 1980). *L. caponis* infestations were diagnosed in April, July, August, November (Islam *et al.*, 1999). In terms of ectoparasites prevalence, chicken breeds were in order: Harco > Lumacco > Rhode Island Red > Barbrock (Ugochukwu & Omije, 1986). The lice are most prevalent in summer seasons i.e. June, July & August and scarcest from November to February (EL-Kifl *et al.*, 1973). The optimum temperature for the development of *Manacanthus stramineus* is 37.7 – 41.5C (Brown, 1970). *Menacanthus pallidulus*, *Menopon gallinae*, *Goniodes dissimilis* and *Goniocotes gallinae* were most abundant on the back region of the host and *Lipeurus caponis* on the wing feathers (Manuel & Anceno, 1981).

**Chemotherapy.** A number of chemicals and herbal products are used for the control of biting and sucking lice infestation of poultry. Malix, a dusting powder containing derris (Hohorst, 1939). Phenol and Cresols at 2% kill lice (Buxton, 1940). Complete control of *M. stramineus* on hens by treating with 100 mg Coumaphos (Co-Ral) dust per square fut (Knapp, 1962). Lindae and BHC (Ware & Naber, 1962). 5% granules of ronnel, 2% granules of bromophos and 2% dust of Dimetilan highly effective against lice infestation (Hoffman & Hogan, 1967). *B.thuringiensis* (Hoffman & Gingrich, 1968). 2.7% butanate and 0.5% lindane supprey (Jungmann *et al.*, 1970). 2% trichlorphon (Puchkova – EA, 1977). Biollethrin, Biopresmetrin, Neo-pinamin and Pybuthrin were effective in low concentration against *Menacanthus stramineus* (Kachekova & Frolov, 1978). Dichlorvos (Nuvan. 7) effective for lice control (Grzywinski *et al.*, 1980). 0.5-1% solution of Phosalone (Zolone) effective against lice infestation (Reddy *et al.*, 1980). Coumaphos (as Asuntol-50) Malathion, Trichorphan (Neguvon) Pflspray, Permethrin (Coopex) and Carbaryl were highly effective in killing lice (Manuel & Macetangay, 1981). The effectiveness of 5% Methyl bromfenvinphos (Polwet. 5) against the fowl ectoparasite (Zlotorzycza *et al.*, 1982). 2% Aqueous solution of Pestoban (Herbal products) very effective against lice infestation (Ahmed, 1986). Pestoban application at a dilution at 1:30 was sufficient to control lice infestation (Sinha *et al.*, 1989). Bacilan, Dipel (a commercial preparation of *Bacillus thuringiensis* subsp. *Kurstaki*) *B.thuringiensis* the serotypes *Kurstaki*, *Finitimus*, *Kenya* and *Morrisoni* were most toxic to *M. gallinae* (Lonc *et al.*, 1988). Permethrin (Pavlovic *et al.*, 1989). Pyrethroids are effective against ectoparasites (Salisch, 1989). Vixon (Carbaryl preparation) highly effective against sucking and biting lice (Werner *et al.*, 1989). The Guinea fowl and chicken were successfully treated with DDT powder and the Peahen with Coumaphos (Okaeme, 1989). Cypermethrin (Chhabra & Donora, 1994). A single treatment with Gammatox as 1% dip was effective in controlling *L. caponis* infestation (Islam *et al.*, 1999).

**Biological control.** Three biological products were available in the USSR for ectoparasite control: two of them (Entobacterin and Dendrobacillin) were preparations of spores and endotoxin of *Bacillus thuringiensis* and the third (Boverin) Conidiospores of the Fungus *Beauveria bassiana*. These were active, alone or in combination with chemical pesticides, against lice infestation (Frolov *et al.*, 1974).

## CONCLUSION

In the light of this review it is concluded that further research is to be conducted into the lice infestation because it is a common problem in poultry which causes heavy in poultry by causing damage to the birds and their production.

## REFERENCES

- Agarwal, G.P. and A.K. Saxena, 1981. The perioesophageal nephrocytes of some ischnoceran Mallophaga (Phthiraptera). *Angewandte Parasitologie*, 22: 104–6
- Ahmad, J., 1986. Use of pestoban for the control of lice in poultry. *Int. J. Indigenous Med.* 5: 11–3
- Abdul Karim, R.M., B.A. Jassim and N.S. Naser, 1988. Infestation and occurrence of ectoparasites (lice, ticks, and mites) of pigeons in Erbil area. *Bull. Ent.* 29: 173–7
- Ahmed, S.M., 1991. Toxonomical study of ectoparasites on indigenous poultry and effect of fowl tick *Argas persicus* on different blood parameters. *M.Sc. Thesis*, College of Veterinary Science, Lahore
- Amin-Babjee, S.M., C.C. Lee and A.A. Mahmood, 1997. Prevalence of ectoparasite infestation in different age groups of village chickens. *J. Vet-Malaysia*. 9: 55–9
- Axtell-RC., 1999. Poultry integrated pest management: status and future. *Integrated-pest-management-reviews*, 4: 53–73
- Buxtan, P.A., 1940. The louse: Present knowledge and future work. *Trans. Royal Soc. Trop. Med. Hyg.*, 33: 365–88
- Buxtan, P.A., 1940. Temperatures lethal to the louse. *Brit. Med. J. Mar.* 2<sup>nd</sup>: 341
- Buxtan, P.A., 1940. The control of lice. *British Med. J.*, Nov. 2<sup>nd</sup>: 603–4
- Brown, N.S., 1970. Distribution of *M. stramineus* in relation to chicken surface temperature. *J. Parasitol.*, 56: 1205
- Buriro, S.N., 1982. Incidence and epizootology of ecto and endo parasites in urban and rural areas of Sind. PARC Research Project. Final Technical Report. Nov. 1980 to Oct. 1981. :57
- Bilqees, F.M. and A. Khan, 1985. Incidence of parasitic infestation in fowl of Karachi. *Pakistan J. Zool.*, 17: 306–8
- Bilinski, S.M. and W. Jankowska, 1987. Oogenesis in the bird louse *M. stramineus* (Insecta, Mallophaga). *Int. J. Zool.*, 116: 1–12
- Cariaso, B.L., 1982. Selected ectoparasites towards efficient management of livestock and poultry. *Philippine Entomol.*, 5: 321–43
- Chandra, S., G.P. Agarwal and A.K. Saxena, 1988. Seasonal changes in the population of Mallophaga. *Angewandteparasitologie*, 29: 244–9
- Chhabra, R.C. and N. Donora, 1994. Ectoparasites of poultry in Zimbabwe and their control. *Zimbn. Vet. J.*, 25: 26–32
- Devi, N.C.T. and P.N. Mishra, 1981. Record of some avian biting lice from Kathmandu Valley. *J. Nat. Hist. Museum*, 5: 11–22
- De-Vaney, J.A., 1986. Ectoparasites in poultry. *Poult. Sci.*, 65: 649–56
- Dikaev, B. Yu., 1988. Ectoparasites of poultry and their control. *Veterinariya, Moscow*, 10: 47–8
- Erllich, I., 1942. Mallophaga of domestic fowl. *Vet. Arhiv.*, 12: 397–406
- Erllich, I., 1942. Biting lice of Turkeys. *Vet. Arhiv.*, 12: 492–4
- Edgar, E.R. and D.F. King. 1950. Effect of the body louse, *Eomenacanthus stramineus*, on mature chickens. *Poult. Sci.*, 29 :214–9
- El-Kifil, A.H., A. Wahab, M.K. Kamel and W.A.E. Abdel, 1973. Poultry ectoparasites in sharia Governorate. *Agri. Rev.*, 51: 113–20
- Frolov, B.A. 1974. Chemical and biological methods of controlling poultry ectoparasites. *Veterinaria – Moscow*, 12: 66–8
- Fabiyi, J.P., 1980. Survey of lice infesting domestic fowls of the Jos Plateau, Northern Nigeria. *Bull. Anim. Hlth. Prod.*, 28: 21–9
- Fedorenko, I.A., N.T. Nikitchenko and V.YA. Kuz'menko, 1983. Bird lice (Mallophaga) of some birds in the central Dnieper area. *Vestnik Zoologii.*, 4: 85–9
- Frolov, B.A., R.A. Li, L.P. Vyunov, V.E. Sivokhina and A.A. Berlim, 1984. Insecticidal and acaricidal activity of cyclophas in relation to ectoparasites of bird. *Vetrinaria-Moscow*, 7: 24–7
- Giles, E.E. and E.S. raun, 1959. Effects of chicken body louse infestation on egg production. *J. Econ. Ent.* 52: 358–9
- Grzywnski, L., M. Mazur Kiewicz and T. Martynowicz, 1980. The effectiveness of Nuvan-7 in the control of external parasites of fowl in intensive production. *Medycyna Weterynaryjna*, 36: 615–8
- George, J.B.D., otobo, J. Ogunleye and B. Adediminyi, 1992. Louse and mites infestation in domestic animals in North Nigeria Trop. *Anim. Hlth. Prod.*, 24: 121–4
- Hoyle, W.L., 1938. Transmission of poultry parasites by birds with special reference to the "English" or House sparrow and chickens. *Trans. Kans. Acad. Sci.*, 41: 379–84
- Hohorst, W., 1939. Mallophaga of hen and their oviposition. *Vet. Med. Nachr.*, 4: 61–88
- Hohorst, W., 1942. Little known Mallophaga of fowls. *Senckenbergiana*, 25: 222–5
- Hathaway, C.R., 1943. Commensalism of biting lice hippoboscid flies on an a pigeon. *Mem. Inst. Osw. Cruz.*, 38: 413–7
- Hoffman, R.A. and B.F. Hogan, 1967. Control of chicken body, shaft and wing lice on laying hens by self treatment with insecticide dusts and gramules. *J. Econ. Ent.*, 60: 1703–5
- Hoffman, R.A. and R.E. Gingrich, 1968. Dust containing *Bacillus thuringiensis* for control of chicken body, shaft and wing lice. *J. Econ. Ent.* 61: 85–8
- Hashmi, F.Q., 1971. A study on the incidence of endo and ecto parasites of wild birds. *M.Sc. Thesis*, West Pakistan Agricultural University, Lyallpur
- Horning, D.S., R.L. Plama, R.L.C. Pilgrim, 1980. The lice (Insecta: Phthiraptera) from the snares Islands, New Zealand Miscellaneous series, National Museum of New Zealand. No. 3, P:17
- Hofstad, M.S., H.J. Barnes, B.W. Calnek and W.M. Yoder., 1984. Diseases of poultry. 8<sup>th</sup> Ed. Iowa State University Press, Ames, Iowa, U.S.A. : 671–700
- Iqbal, M., 1971. Studies on the ectoparasites of the livestock with special emphasis on the incidence, economic losses and chemotherapy. *M.Sc. Thesis*, West Pakistan Agricultural University, Lyallpur
- Iqbal, M., M. Ajmal, A.H. Choudhry and A.R. Kazam. 1971. Studies on the ectoparasites infesting experimental livestock and poultry of West Pakistan Agricultural University, Lyallpur. Proc. 23 rd Pak. Sci. Conf., J14
- Islam, M.K., M.M.H. Mondal, M.M. Rehman, A.K.M.F. Haque and M.A.A. Chaudhery, 1999. Effects of *Lipeurus caponis* Linnaeus, 1958 (Mallophaga: Philopteridae) on laying hens. *Vet. Rev.*, 14: 32–3
- Jungmann, R., R. Ribbeck, S. Eisen blatter and H. Schematus, 1970. Infestation of laying hens with *Dermanyssus gallinae* and feather lice: Harmful effects and control. *Mh. Ver. Med.*, 25: 28–32
- Knapp, F.W., 1962. Co-Ral as a litter and nest dust to control the chicken body louse. *J. Econ. Ent.*, 55: 571–2
- Kumar, K. and B.N. Sahai, 1974. On the incidence of poultry lice in desi fowls. *Indian J. Anim. Hlth.*, 13: 165
- Kachekova, Sh. and B.A. Frolov, 1978. Insecticidal and acaricidal activity of pyrethroids against poultry ectoparasites. *Problems- veterinarnoi-Sanitarii*, 61: 61–3
- Kaminjolo, J.S., E.S. Tibasingh and G.A. Ferdinand, 1988. Parasites of the common pigeon (*Columba livie*) from the environs of port of spain, Trinidad. *Bull. Anim. Hlth. Prod. Africa*, 36: 194–5
- Lonc, E., T.M. Lachowicz and M. Mazurviewixz, 1988. Efficacy of *Bacillus thuringiensis* against poultry parasites. *Zeszyt Naukowe Akademick Rolniczej wroclawiu, weterynaria*, 45: 63–6
- Larramendy, R., B.N. Szczypel, A. Perez, A. Gonzalez and J. Estrada, 2000. Efficacy of neem oil (*Azadirachta indica* A, Juss) in hens naturally

- infestated with ectoparasites. *Revista Cubana-de-Ciencia-Avicola*, 24: 125–31
- Mathur, R.P., 1973. Control of ectoparasites during the rainy season. *Poult. Guide*, 22–31
- Martin, M.P., 1973. Spanish species of Meno–ponidae (Mallophaga) parasitic on poultry. *Revista Iberica de parasitologica*, 33: 281–94
- Martin–Mateo, M.P., F. Albalan and C. Sanchez Acedo, 1980. Instituto Espanol de Entomologia. *Calle J. Gutierrez Abascal*, 34: 121–45
- Manuel, M.F., 1981. The ectoparasite (lice and mites) occurring in domestic chickens in the Philippines. *J. Vet. Med.* 20: 87–100
- Manuel, M.F. and V.C. Macatangay, 1981. Efficacy trials of some insecticidal preparations against common lice and mites effecting domestic chicken in Philippines. *J. Vet. Med.* 20: 58–70
- Manuel, M.F. and T.A. Anceno, 1981. Distribution of biting lice (Mallophaga) on the body of netive chickens (*Gallus gallus domesticus*). *J. Vet. Med.* 20: 50–57
- Mehl, R., 1981. Our knowledge of the fauna of biting lice (Mallophaga) in Norway. *Norge. Fauna, Norway*, 34: 80–5
- Modrzejewska, M., 1982. Mallophaga on pheasants in lower Silesia. *Wiadomosci Parazytologiczne*, 28: 175–6
- Morrow, C., 1986. *Poultry Parasites*. In: 2<sup>nd</sup> Poultry Health Proceeding of the Australian Veterinary Poultry Association, 26–30 May Sydney, Australia
- Noh, Y.T., K.M. Back and I.H. Moon, 1989. Taxonomic studies on chewing lice (Mallophaga, Philopteridae) in Korea. *Korean L. Ent.*, 19: 143–7
- Okaeme, A.N., 1989. Lameness associated with ectoparasitic infestation in *Numidia meleagris galeata*, *Gallus domestica*, *Pavo multicus* (Sic). *Bulletin Anim. Hlth. Prod.*, 37 :189–90
- Phillip, C.B., 1963. In Naegele. J.A. (ed), *Advances in Acarology*, 1:285–325. *Cornel Univ. Press*, Ithaca, N.Y.
- Puchkova–EA., 1977. Elimination of ectoparasites of poultry on poultry farms in USSR. *Veterinariya, Moscow, –USSR*. 7: 19–22
- Panda, D.N. and S.S. Akluwali, 1983. Effects of *M. stramineus* and *L. tropicalis* infestation on weight gain in broiler birds. *Indian Vet. J.*, 60: 85–7
- Pavlovic, I., V. Blazin, V. Hudnia, Z. Iliac and B. Miljkovic, 1989. Effects of the biting louse *M. stramineus* on reducing the egg production of poultry under intensive condidtion. *Veterinarski Glasnik*. 43: 181–6
- Reddy, K.R., J. Rao and K. Rao, 1980. Field trial of “zolone” as an insecticide on poultry lice and bed bugs. *Pesticidas*, 14: 36–7
- Rodriguez Caabeiro, F., A. Jimenez Gonzalez, and S.Hernandez Rodriguez, 1981. *Goniodes pavonis* (Linneo, 1758) parasite of the peacock. *Revista Iberica de Peresitologia.*, 41; 601–3
- Ratanasethakul, C., S. Pholpark, B. Laopaiboon, M. Polpark and D. Tuntasuvan, 1985. Parasites of native chickens in Northeast Thailand. *J. Vet. Med.* 15: 229–42
- Reddy, A.R.M., R. Shudhakar, P.K. Reddy and P.R. Reddy, 1985. Lousiness in poultry. *Poult. Adviser*, 18: 27–9
- Ribbeck, R., G. Heider and G. Manreal, 1992. Arthropod infestation of poultry. *Krankheiten–des–wirtschaftsgeflugles*, 3: 439–64
- Rafiq, N.Z., 2000. Taxonomy and effects of ectoparasites on the egg production and blood parameters of White Leghorn and indigenous layers. *M.Sc. Thesis*, College of Veterinary Science, Lahore
- Sinha, R.P. R.S. Prasad, S. Roy and M. Zahiruddin, 1989. Effect of pestoban against ectoparasites of livestock and poultry. *Livestock Advisor*, 12: 26–30
- Salisch, H., 1989. Recent developments in the chemotherapy of parasitic infections of poultry. *World's Poult. Sci. J.*, 45: 115–24
- Saxena, A.K., A. Kumar and S.K. Singh, 1995. Prevalence of *Menopon gallinae* Linne (Phthiraptera: Amblycera) on poultry birds of Garhwal. *J. Parasit. Dis.*, 19: 69–72
- Shafique, M., 1979. A study on the ectoparasites of commercialized poultry in Punjab. *M.Sc. Thesis*, University of Agriculture, Faisalabad
- Stockdale, H.J. and E.S. Raun, 1960. Economic importance of chicken body louse. *J. Econ. Ent.*, 53; 421–3
- Singh, A. and R.C. Chhabra. 1973. Incidence of arthropod pests of domestic animals and birds. *Indian J. Anim. Sci.*, 43: 393–7
- Soulsby, E.J.L., 1982. *Helminths, Arthropods and Protozoa of Domesticated Animals*, 7<sup>th</sup> Ed., pp: 366–87. Baillere Tindall and Cassell Ltd., London
- Shavsmanou, S.H., 1982. A contribution to a study of the bird lice (Mallophaga) of domestic and wild birds of the Dzhizak reition. *Doklady Akademii Nauk Uzbekshii SSR.*, 4: 50–1
- Shahjahan, I.A., and S. Iqbal, 1995. Prevalence of chickens’s lice (Mallophaga) in Peshawar and Mansehra (NWFP). *Pakistan Vet. J.*, 15: 58–60
- Tigin, Y., 1973. Ectoparasites of domestic pigeons (*columba livia*) in Turkey. *Veteriner Fakultesi Dergisi*, 20: 372–90
- Urban, E., and J. Zlotorzycza, 1981. Investigation on the vitality of farul Mallophaga after the loss of their host. *Angewandte parasitologie*, 22: 83–91
- Ugochukwu, E. and F.A. Omije, 1986. Ectoparasitic fauna of poultry in Nsukka. *Int. J. Zoonoses*, 13: 93–7
- Urquhart, G.M., 1987. *Veterinary Parasitology*, 1<sup>st</sup> Ed., pp: 256–7. ELBS, Longman House, Burnt Mill, Harlow, England
- Ware, G.W. and E.C. Naber, 1962. Lindene and BHC in egg yolks following recommended uses for louse and mite control. *J. Econ. Ent.*, 55: 568–70
- Werner, G. Hiepe, T.Tschavschev, P. Hahn, H. Zimmermann, H. Siering and L.W. Hoffmann, 1989. Vixon powder a carbayl preparation against ectoparasites in domestic animal. *Monatshefte–for–veterinarmedizin*, 44: 611–3
- Zlotorzycza, J., 1980. Ectoparasites of synantropic birds and mammals. *Wiadomosci parazytologiczne*, 28: 157–62
- Zlotorzycza, J., Lonc, E. Giebel, O. Mazurkiewicz and Z.M. Wachnik, 1982. Susceptibility of ectoparasites of fowls to the activity of methyl bromfenvinphos (Polwet. 5). *Wiadomosci parazytologiczne*, 28: 387–392
- Zhang–Deling, Jia–Jnn Yuan, Zhang–Goosheng, DL. Zhang, , Jia–JY and Zhang–GS., 1996. An investigating of poultry diseases in Gansu. *Chinese J. Vet. Med.*, 22: 6–27
- Mathysse, J.G., 1972. *External Parasites*. In: *Diseases of Poultry*, pp: 793–831. Hofstad Press, Ames, Iowa, U.S.A

(Received 01 December 2002; Accepted 10 March 2003)