

## Research Article

# Knowledge, Attitude, and Practice of *Pediculus Capitis* Prevention and Control and Their Predictors among Schoolchildren in Woreta Town, Northwest Ethiopia, 2018: A School-Based Cross-Sectional Study

Henok Dagne <sup>1</sup>, Awel Aba Biya,<sup>1</sup> Amanuel Tirfie,<sup>1</sup> Walelegn Worku Yallew <sup>1</sup>,  
Zewudu Andualem <sup>1</sup>, and Baye Dagne <sup>2</sup>

<sup>1</sup>Department of Environmental and Occupational Health and Safety, Institute of Public Health, College of Medicine and Health Sciences, University of Gondar (UoG), P.O. Box 196, Gondar, Ethiopia

<sup>2</sup>Department of Human Physiology, School of Medicine, UoG, P.O. Box 196, Gondar, Ethiopia

Correspondence should be addressed to Henok Dagne; enoch2313@gmail.com and Baye Dagne; bayedagne7@gmail.com

Received 18 May 2020; Revised 2 June 2020; Accepted 8 June 2020; Published 1 July 2020

Academic Editor: Frederick J. Kaskel

Copyright © 2020 Henok Dagne et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Background.** *Pediculus capitis* is a human head lice infestation, a major public health issue that is most prevalent in resource-limited countries globally. The current study aimed to assess the knowledge, attitude, and practice of *pediculus capitis* prevention and control and their predictors among schoolchildren in North West Ethiopia. **Methods.** About 402 randomly selected schoolchildren from three schools in Woreta town participated in the study from April to June 2018. The outcomes of this study were knowledge, attitude, and self-reported practice of schoolchildren about *pediculus capitis* prevention and control. We used EPI Info 7.1 and SPSS 21 software for data entry and analysis, respectively. Binary logistic regression was employed to test the association of covariates with the outcome/response variables. Variables with a  $p$  value  $<0.2$  during the bivariable binary logistic regression analysis were included in the multivariable binary logistic regression analysis. Variables with  $p$  value  $<0.05$  were declared as significantly associated with outcomes. **Results.** The mean age of the study participants was 10.19 ( $\pm 1.62$ ) years. About 58.8%, 45.8%, and 78.6% of the schoolchildren had better self-reported *pediculus capitis* prevention knowledge, attitude, and practice, respectively. Age of children [9 to 11 years (AOR = 2.24, 95% C.I (1.10, 4.55)) and  $>12$  years (AOR = 3.84, 95% C.I (1.56, 9.46))], better practice (AOR = 2.93, 95% C.I (1.39, 6.18)), and those who were not infested (AOR = 2.25, 95% C.I (1.14, 4.44)) were predictors of knowledge regarding *pediculus capitis* prevention. Better practice (AOR = 4.33, 95% C.I (1.69, 11.09)) and absence of infestation (AOR = 2.97, 95% C.I (1.64, 5.36)) were predictors of attitude of schoolchildren about *pediculus capitis* prevention. Number of students in a class [51 to 56 students per classroom, AOR = 4.61, 95% C.I (1.83, 11.67); 57 to 58 students per classroom, AOR = 8.18, 95% C.I (2.73, 24.46)], less than five family size (AOR = 2.37, 95% C.I (1.24, 4.54)), better knowledge (AOR = 2.93, 95% C.I (1.32, 6.50)), desirable attitude (AOR = 4.24, 95% C.I (1.60, 11.23)), and absence of infestation (AOR = 3.52, 95% C.I (1.22, 10.15)) were predictors of self-reported *pediculus capitis* prevention practice. **Conclusion.** The knowledge, attitude, and practice of schoolchildren regarding *pediculus capitis* prevention and control were not satisfactory. To bring change, intensive efforts on factors associated with the knowledge, attitude, and practice should be encouraged.

## 1. Background


*Pediculus capitis* is infestation by head lice that is a major public health problem globally which is most prevalent in low-income countries [1]. It is an omnipresent issue in children [2] in both schools and in the broader community [3].

*Pediculus capitis* can cause loss of sleep, irritation, pruritus, discomfort, secondary bacterial infections (such as impétigo and acute glomerulonephritis), and lymphadenopathy [4, 5]. The head lice are a potential vector for disease-causing organisms such as *Rickettsia prowazekii*, *Bartonella quintana*, and *Borrelia recurrentis* [6–11]. The methods to reduce the

TABLE 1: Explanatory variables of school children Woreta town, 2018 ( $n = 402$ ).

Variables	Categories	Frequency	Percent (%)
Student grade level	Grade1	99	24.6
	Grade2	99	24.6
	Grade3	100	24.9
	Grade4	104	25.9
Sex	Male	186	46.3
	Female	216	53.7
Age	5-8	85	21.1
	9-11	230	57.2
	>12	87	21.6
Religion	Orthodox	291	72.4
	Muslim	108	26.9
	Protestant	3	0.7
Fathers education	Illiterate	111	27.6
	Elementary	161	40
	Secondary and above	130	32.3
Mothers education	Illiterate	171	42.5
	Elementary	162	40.3
	Secondary and above	69	17.2
Fathers occupation	Private worker	153	38.1
	Government worker	64	15.9
	Daily labor	42	10.4
	Others	143	35.6
Mother's occupation	Private worker	62	15.4
	Government worker	25	6.2
	Housewife	240	59.7
	Others	75	18.7
Family size	≤5	227	56.4
	>5	175	45.6
Personal hygiene control	Yes	106	26.4
	No	296	73.6
Water access	Yes	28	7
	No	374	93
Average number of students per classroom	40-50	116	28.9
	51-56	121	30.1
	57-58	89	22.1
	59-67	76	18.9

morbidity and prevalence of *pediculus capitis* include strategies targeted at increasing knowledge, changing attitudes and behaviors, and improving personal hygiene practice [6]. Even though *pediculus capitis* is very common in Africa from earlier times until now [12–15], there are only a few studies on knowledge, attitude, and practice regarding head lice infesta-

FIGURE 1: Knowledge, attitude, and practice towards *Pediculus capitis* prevention and control among schoolchildren Woreta, 2018 ( $n = 402$ ).TABLE 2: Correlation between knowledge, attitude, and practice towards *Pediculus capitis* prevention and control among schoolchildren Woreta, 2018 ( $n = 402$ ).

Variables	Correlation coefficient	$p$ value*
Knowledge-attitude	0.629	<0.001
Knowledge-practice	0.363	<0.001
Attitude-practice	0.370	<0.001

\*Correlation significant at  $p < 0.01$ .

tions especially in sub-Saharan Africa [16]. Above all, majority of the previous studies focused on the knowledge, attitude, and practice of parents, nurses, and teachers only [17–21]. Let alone the level of knowledge about human head lice and their control in high-income countries is restricted in resource-constrained settings among schoolchildren [16, 19, 20, 22] and even insufficient among health professionals [23–25]. The major setbacks for ineffective control of *pediculus capitis* include lack of knowledge, undesirable attitude towards control and prevention of head lice, and inadequate personal hygiene practice [26–28].

Therefore, the present study was undertaken to assess the level of knowledge, attitude, practice, and their associated factors regarding *pediculus capitis* prevention and control among schoolchildren in Woreta town, northwest Ethiopia.

## 2. Methods

**2.1. Study Design and Setting.** This was a school-based cross-sectional study conducted from April to June 2018 among schoolchildren in Woreta town primary schools. The town is located 589 km far from Addis Ababa, the capital city of Ethiopia. During the study period, there were three primary schools in the town with a total of 3239 students. The prevalence of *pediculus capitis* during the study period was 65.7% (95% CI 60.01–70.3%) [29].

TABLE 3: Factors associated with knowledge regarding *pediculus capitis* prevention and control among schoolchildren Woreta, 2018 ( $n = 402$ ).

Variables	Categories	Knowledge		COR (95% CI)	AOR (95% CI)
		Good (%)	Poor (%)		
Age in years	5 to 8	37 (43.5)	48 (56.5)	1	1
	9 to 11	136 (59.1)	94 (40.9)	1.88 (1.14, 3.10)	2.24 (1.10, 4.55)*
	≥12	63 (72.4)	24 (27.6)	3.41 (1.80, 6.43)	3.84 (1.56, 9.46)**
Fathers education	Illiterate	53 (47.7)	58 (52.3)	1	1
	Elementary	93 (57.8)	68 (42.2)	1.50 (0.92, 2.43)	1.18 (0.55, 2.56)
	Secondary and above	90 (69.2)	40 (30.8)	2.46 (1.45, 4.17)	1.73 (0.65, 4.64)
Mothers education	Illiterate	88 (51.5)	83 (48.5)	1	1
	Elementary	99 (61.1)	63 (38.9)	1.48 (0.96, 2.29)	0.80 (0.37, 1.72)
	Secondary and above	49 (71)	20 (29)	2.31 (1.27, 4.21)	0.73 (0.24, 2.22)
Number of students per class	40-50	78 (67.2)	38 (32.8)	2.03 (1.12, 3.67)	1.77 (0.76, 4.16)
	51-56	62 (51.2)	59 (48.8)	1.05 (0.59, 1.86)	0.68 (0.29, 1.61)
	57-58	58 (65.2)	31 (34.8)	1.87 (1.00, 3.50)	0.96 (0.38, 2.45)
	59-67	38 (50)	38 (50)	1	1
Family size	≤5	146 (64.3)	81 (35.7)	1.70 (1.14, 2.55)	1.00 (0.56, 1.80)
	>5	90 (51.4)	85 (48.6)	1	1
Water accessibility	Yes	13 (46.4)	15 (53.6)	0.59(0.27, 1.27)	0.75 (0.26, 2.16)
	No	223 (59.6)	151 (40.4)	1	1
Attitude towards pediculus capitis prevention	Poor	66 (30.3)	152 (69.7)	1	1
	Good	170 (92.4)	14 (7.6)	27.96(15.09, 51.82)	16.95 (8.74, 32.88)***
Practice towards <i>pediculus capitis</i> prevention	Poor	21 (24.4)	65 (75.6)	1	1
	Good	215 (68)	101 (32)	6.59 (3.82, 11.37)	2.93 (1.39, 6.18)**
<i>Pediculus capitis</i> infestation	Yes	116 (84.1)	22 (15.9)	1	1
	No	120 (45.5)	144 (54.5)	6.33 (3.78, 10.6)	2.25 (1.14, 4.44)*

Significant at \* $p \leq 0.05$ , \*\* $p \leq 0.01$ , \*\*\* $p \leq 0.001$ , Hosmer-Lemeshow goodness-of-fit ( $p = 0.196$ ) 1 reference.

**2.2. Sample Size Calculation and Sampling Procedure.** For the present study, the sample size was determined using a single population proportion formula [30]. With the following assumptions:  $p = 50$  percent to allow maximum variation (as there was no previous country analysis on the ratio of knowledge, attitude, and practice to *pediculus capitis* prevention and control), 95 percent confidence level,  $z =$  standard normal tabulated value, and  $\alpha =$  level of significance and margin of error ( $d$ ) = 0.05;

$$n = \frac{(z_{\alpha/2})^2 p(1-p)}{d^2} = \frac{(1.96)^2 0.5(1-0.5)}{0.05^2} = 384 \quad (1)$$

The final total sample size was 402, after adding the estimated nonresponse rate of 5 percent. Study participants were chosen using a simple random sampling technique and distributed in proportion to the three schools, based on the number of students in each school.

**2.3. Data Collection Tool and Quality Control Procedures.** A pretested, semistructured questionnaire was used that included socio-demographic variables, knowledge, attitude, and practice items relevant to infestation with *pediculus capitis*. Two Environmental Health Bachelor's degree students

conducted an interview and observation after receiving training on the data collection method, techniques, study intent, and ethical considerations.

Detailed explanation and reliability of the method for data collection based on the pretest results are discussed elsewhere [29].

#### 2.4. Variable Measurement

**2.4.1. Pediculosis.** A child with at least one head louse by wet combing is assumed to be infested with *pediculus capitis* [14].

**2.4.2. Schoolchildren.** In the current report, children attending classes from grades 1 to 4 were considered schoolchildren [29].

**2.4.3. Knowledge.** Knowledge was measured by 10 yes/no category knowledge items.

Students who scored mean of knowledge questions and above were deemed to have good knowledge [29].

**2.4.4. Attitude.** Attitude was assessed by 8 Likert-scale attitude questions (1-strongly disagree with 5-strongly agree). Children who scored mean and above questions about the attitude were known as having a good attitude [29].

TABLE 4: Factors associated with attitude regarding *pediculus capitis* prevention among schoolchildren Woreta, 2018 ( $n = 402$ ).

Variables	Categories	Attitude		COR (95% CI)	AOR (95% CI)
		Good (%)	Poor		
Age in years	5 to 8	30 (35.3)	55 (64.7)	1	1
	9 to 11	105 (45.7)	125 (54.3)	1.54 (0.92, 2.58)	1.18 (0.56, 2.46)
	≥12	49 (56.3)	38 (43.7)	2.36 (1.28, 4.37)	1.47 (0.61, 3.57)
Fathers education	Illiterate	35 (31.5)	76 (68.5)	1	1
	Elementary	76 (47.2)	85 (52.8)	1.94 (1.17, 3.22)	1.87 (0.84, 4.17)
	≥secondary	73 (56.2)	57 (43.8)	2.78 (1.64, 4.71)	1.65 (0.61, 4.44)
Mothers education	Illiterate	64 (37.4)	107 (62.6)	1	1
	Elementary	78 (48.1)	84 (51.9)	1.55 (1.00, 2.40)	0.72 (0.34, 1.53)
	≥secondary	42 (60.9)	27 (39.1)	2.6 (1.46, 4.62)	0.90 (0.38, 2.20)
Number of students per class	40-50	56 (48.3)	60 (51.7)	1.49 (0.82, 2.68)	0.78 (0.34, 1.81)
	51-56	51 (42.1)	70 (57.9)	1.18 (0.66, 2.12)	0.92 (0.38, 2.20)
	57-58	48 (53.9)	41 (46.1)	1.90 (1.02, 3.54)	0.96 (0.39, 2.40)
	59-67	29 (38.2)	47 (61.8)	1	1
Family size	≤5	119 (52.4)	108 (47.6)	1.86 (1.25, 2.79)	1.19 (0.67, 2.12)
	>5	65 (37.1)	110 (62.9)	1	1
Knowledge towards <i>pediculus capitis</i> prevention	Poor	14 (8.4)	152 (91.6)	1	1
	Good	170 (72)	66 (28)	27.96 (15.09, 51.82)	17.32 (8.90, 33.72)***
Practice towards <i>pediculus capitis</i> prevention	Poor	9 (10.5)	77 (89.5)	1	1
	Good	175 (55.4)	141 (44.6)	10.62 (5.14, 21.93)	4.33 (1.69, 11.09)**
<i>pediculus capitis</i> infestation	Yes	81 (30.7)	183 (69.3)	1	1
	No	103 (74.6)	35 (25.4)	6.65 (4.18, 10.58)	2.97 (1.64, 5.36)***

Significant at  $*p \leq 0.05$ ,  $**p \leq 0.01$ ,  $***p \leq 0.001$ , Hosmer-Lemeshow goodness-of-fit ( $p = 0.134$ ). COR: crude odds ratio; AOR: adjusted odds ratio; CI: confidence interval.

**2.4.5. Practice.** Children were asked five specific questions regarding the prevention behavior of *pediculus capitis*. Those children who scored mean and above of the questions were considered good practice [29].

**2.5. Data Analysis.** Completeness and consistency of the data were reviewed regularly. Epi info version 7.1 was used for data entry and analysis was performed using SPSS version 21. We calculated the number, mean, and standard deviation to show the descriptive results.

Bivariable binary logistic regression was first tested, and in the final model, we used multivariable binary logistic regression to test variables with a  $p$  value  $< 0.2$  in bivariable analysis for significant association. Variables with a  $p$  value  $< 0.05$  were declared as being significantly associated with dependent variables (i.e., knowledge, attitude, and practice) in the multivariable binary logistic regression analysis.

Crude and adjusted odds ratios and 95% confidence intervals were calculated. Model fitness was tested using Hosmer and Lemeshow goodness of fit at  $p > 0.05$ .

**2.6. Patient and Public Participation.** The current research did not include patients or the public but included students with and without *pediculus capitis* infestation.

### 3. Results

**3.1. Socio-Demographic Characteristics.** Four hundred and two students were included in the current study. Above half (53.7%) of the study participants were females. The study participants' mean age was 10.19 ( $\pm 1.62$ ) years. Almost all (93%) of the students attend in schools with no water access (Table 1).

From the total participants, 78.6% had good self-reported practice, 58.8% had good knowledge, and 45.8% had desirable attitude (Figure 1).

**3.2. Correlation between Knowledge, Attitude, and Practice.** The correlation between knowledge, attitude, and practice was assessed using the rank correlation coefficient of Spearman and represented using  $0 - 0.25 =$  poor,  $0.25 - 0.5 =$  average,  $0.5 - 0.75 =$  good, and above  $0.75 =$  excellent correlations based on criteria developed for the study of statistical power for behavioral sciences [31]. There was a significant correlation among knowledge, attitude and practice (Table 2).

**3.3. Factors Associated with Knowledge regarding *pediculus capitis* Prevention.** Age of child, paternal and maternal education, number of students per class, family size, water accessibility, attitude of students towards *pediculus capitis* prevention, practice towards *pediculus capitis* prevention,

TABLE 5: Factors associated with *pediculus capitis* prevention practice among schoolchildren Woreta, 2018 ( $n = 402$ ).

Variables	Categories	Practice		COR (95% CI)	AOR (95% CI)
		Good (%)	Poor (%)		
Grade level	Grade 1	76 (76.8)	23 (23.2)	1	1
	Grade 2	66 (66.7)	33 (33.3)	0.61 (0.32, 1.13)	0.32 (0.13, 0.76)
	Grade 3	79 (79)	21(21)	1.14 (0.58, 2.23)	0.62 (0.25, 1.52)
	Grade 4	95 (91.3)	9 (8.7)	3.19 (1.40, 7.31)	0.57 (0.18, 1.80)
Fathers education	Illiterate	72 (64.9)	39 (35.1)	1	1
	Elementary	125 (77.6)	36 (22.4)	1.88 (1.10, 3.22)	0.92 (0.43, 1.98)
	Secondary and above	119 (91.5)	11 (8.5)	5.86 (2.82, 12.16)	1.46 (0.47, 4.50)
Mothers education	Illiterate	113 (66.1)	58 (33.9)	1	1
	Elementary	137 (84.6)	25 (15.4)	2.81 (1.65, 4.78)	1.29 (0.56, 2.92)
	≥secondary	66 (95.7)	3 (4.3)	11.29 (3.40, 37.47)	5.36 (0.98, 29.36)
Number of students per class	40-50	92 (79.3)	24 (20.7)	2.94 (1.55, 5.57)	2.20 (0.90, 5.38)
	51-56	99 (81.8)	22 (18.2)	3.45 (1.81, 6.60)	4.61 (1.83, 11.67)***
	57-58	82 (92.1)	7 (7.9)	8.99 (3.67, 22.01)	8.18 (2.73, 24.46)***
	59-67	43 (36.6)	33 (43.4)	1	1
Family size	≤5	197 (86.8)	30 (13.2)	3.09 (1.88, 5.09)	2.37 (1.24, 4.54)**
	>5	119 (68)	56 (32)	1	1
Knowledge towards <i>pediculus capitis</i> prevention	Poor	101 (60.8)	65 (39.2)	1	1
	Good	215 (91.1)	21 (8.9)	6.56 (3.82, 11.37)	2.93 (1.32, 6.50)**
Attitude towards <i>pediculus capitis</i> prevention	Poor	141 (64.7)	77 (35.3)	1	1
	Good	175 (95.1)	9 (4.9)	10.62 (5.14,21.93)	4.24 (1.60, 11.23)**
<i>Pediculus capitis</i> infestation	Yes	184 (69.7)	80 (30.3)	1	1
	No	132 (95.7)	6 (4.3)	9.56 (4.05, 22.57)	3.52 (1.22, 10.15)*

Significant at \* $p \leq 0.05$ , \*\* $p \leq 0.01$ , \*\*\* $p \leq 0.001$ , Hosmer-Lemeshow goodness-of-fit ( $p = 0.478$ ) 1 reference.

and previous history of *pediculus capitis* infestation were variables candidate for the multivariable binary logistic regression analysis as they have  $p$  value less than 0.2. Only age, attitude towards *pediculus capitis* prevention, practice towards *pediculus capitis* prevention, and *pediculus capitis* infestation were associated with knowledge regarding *pediculus capitis* prevention during the multivariable binary logistic regression analysis. Children aged 9 to 11 years had 2.24 times [AOR = 2.24, 95% C.I (1.10, 4.55)], and those aged  $\geq 12$  years had 3.84 times [AOR = 3.84, 95% C.I (1.56, 9.46)] better *pediculus capitis* prevention knowledge than those aged 5 to 8 years. Schoolchildren with good *pediculus capitis* prevention practice had 2.93 times better knowledge as compared to those with poor practice [AOR = 2.93, 95% C.I (1.39, 6.18)]. Study participants with previous history of *pediculus capitis* had twice better preventive knowledge than those with no history of infestation [AOR = 2.25, 95% C.I (1.14, 4.44)] (Table 3).

Students' knowledge towards *pediculus capitis* prevention, practice towards *pediculus capitis* prevention, and previous history of *pediculus capitis* infestation were factors associated with attitude about *pediculus capitis* prevention during the multivariable logistic regression (Table 4).

Grade level, paternal and maternal educational status, number of students per classroom, family size, knowledge

and attitude towards *pediculus capitis* prevention, and history of infestation were factors eligible ( $p < 0.2$ ) for multivariable analysis in the final model. The only number of students per classroom, family size, knowledge, attitude, and history of infestation was associated with *pediculus capitis* prevention practice during the multivariable analysis. Class size was associated with practice regarding *pediculus capitis* prevention. Students attending in larger size class had better prevention practice compared with those in smaller size classes. Students from lower family size had better *pediculus capitis* prevention practice [AOR: 2.37, 95% C.I (1.24, 4.54)]. Children with good knowledge had 2.93 times better practice than those with poor knowledge [AOR: 2.93, 95% C.I (1.32, 6.50)]. Study participants with desirable attitudes had 4.24 times better prevention practice than those with undesirable attitudes [AOR: 4.24, 95% C.I (1.60, 11.23)]. Children with previous history of infestation by *pediculus capitis* had better prevention practice [AOR: 3.52, 95% C.I (1.22, 10.15)] (Table 5).

#### 4. Discussion

In this study, we evaluated the level of knowledge, attitude, and practice regarding *pediculus capitis* prevention and control among schoolchildren in Woreta town. This study is a



part of a project on a school health assessment program, and detailed information about prevalence and risk factors of *pediculus capitis* is published elsewhere [29]. In the current manuscript, we have assessed the knowledge, attitude, and practice regarding *pediculus capitis* prevention and control and their associated factors. Knowledge was a significant predictor of infestation status in previous studies [19, 29, 32]. This may be due to the lack of knowledge that can result in insufficient capacity to handle lice infestation. Several interventional studies concentrated on health education intervention to improve knowledge of school teachers, guardians, and students [33, 34]. Deficiencies in knowledge may indicate inability to manage infestation. In the current analysis, the positive associations between knowledge-attitude, knowledge-practice, and attitude-practice notify the relationship among knowledge, attitude, and practice about *pediculus capitis* prevention. Sufficient knowledge will result in a positive attitude leading to better practice. This is in line with previous studies [35, 36]. Students with higher age, better attitude and practice, and those with no previous history of infestation had better-adjusted odds of knowledge regarding *pediculus capitis* prevention. Students with higher age were less likely to be infested [5]. In the current study, students with higher age were found to have better infestation prevention and control knowledge. This is in line with earlier studies [37].

Family size less than five, better knowledge and desirable attitude, and not being infested by *pediculus capitis* were among factors associated with practice regarding *pediculus capitis* prevention among schoolchildren. Previous research showed that children from lower socioeconomic classes and those with lower-educated parents were more frequently infested [38–43]. Greater family size was identified as a determinant for *pediculus capitis* in several earlier studies [44–46].

In the current study, students with better knowledge and desirable attitude more likely reported better *pediculus capitis* prevention and control practice. Health practice is defined by the knowledge and attitude of an individual or the public [37].

This research was, ultimately, not without limitations. No evaluation was made of the knowledge, attitude, and practice of school teachers and parents. The scarcity of previous studies on knowledge, attitude, practice, and associated factors among children made comparison of results difficult. An additional limitation of this study has been the inherent weakness of cross-sectional research design in determining cause-effect relationship, recall and social desirability bias, and poor generalizability as the analysis is performed only at a specific city.

## 5. Conclusion

Head lice infestation is a major public health concern and the national and regional health authorities need to advocate awareness-raising programs that target mothers and prepare knowledge, attitude, and practice improvement strategies.

## Abbreviations

AOR: Adjusted odds ratio  
COR: Crude odds ratio

CI: Confidence interval  
EPI Info: Epidemiological information  
SPSS: Statistical Package for Social Sciences.

## Data Availability

The datasets available from the corresponding authors upon reasonable request.

## Ethical Approval

Ethical approval was secured from the department of Environmental and Occupational Health and Safety Ethical committee. Written consent was obtained from parents and school directors. Assent was received from study participants. Any possible identifiers were avoided to ensure confidentiality. Health education about *pediculus capitis* prevention and control, transmission routes, and personal hygiene was delivered for school teachers, students, and directors after completion of data collection. School directors were advised to regularly check the personal hygiene of students. Students with *pediculus capitis* were advised for visiting nearby clinics.

## Conflicts of Interest

None of the authors declared competing interest.

## Authors' Contributions

HD designed and supervised the study, analysed the data, and wrote the draft manuscript. WWY participated in the design of the study, revised the draft manuscript, and critically revised the draft manuscript for the methodological and intellectual soundness. AT, AAB, ZA, and BD involved from proposal writing, participated in data collection, until the manuscript writing phase. The final manuscript was read and accepted by all contributors.

## Acknowledgments

The authors are grateful to the University of Gondar, data collectors, supervisors, Woreta education office, and study participants for their invaluable support and cooperation to conduct the research.

## References

- [1] D. Pilger, J. Heukelbach, A. Khakban, F. A. Oliveira, G. Fengler, and H. Feldmeier, "Household-wide ivermectin treatment for head lice in an impoverished community: randomized observer-blinded controlled trial," *Bulletin of the World Health Organization*, vol. 88, no. 2, pp. 90–96, 2010.
- [2] D. L. Bonilla, L. A. Durden, M. E. Eremeeva, and G. A. Dasch, "The biology and taxonomy of head and body lice—implications for louse-borne disease prevention," *PLoS Pathogens*, vol. 9, no. 11, article e1003724, 2013.
- [3] P. Magalhães, E. V. Figueiredo, and D. P. Capingana, "Head lice among primary school children in Viana, Angola: prevalence and relevant Teachers' knowledge," *Human Parasitic Diseases*, vol. 3, pp. 11–18, 2011.

- [4] A. Doroodgar, F. Sadr, M. Doroodgar, M. Doroodgar, and M. Sayyah, "Examining the prevalence rate of *Pediculus capitis* infestation according to sex and social factors in primary school children," *Asian Pacific journal of Tropical Disease*, vol. 4, no. 1, pp. 25–29, 2014.
- [5] A. Vahabi, K. Shemshad, M. Sayyadi et al., "Prevalence and risk factors of *Pediculus (humanus) capitis* (Anoplura: Pediculidae), in primary schools in Sanandaj City, Kurdistan Province, Iran," *Tropical Biomedicine*, vol. 29, no. 2, pp. 207–211, 2012.
- [6] M. Yingklang, C. Sengthong, O. Haonon et al., "Effect of a health education program on reduction of pediculosis in school girls at Amphoe Muang, Khon Kaen Province, Thailand," *PLoS One*, vol. 13, no. 6, article e0198599, 2018.
- [7] N. Amanzougaghene, F. Fenollar, A. K. Sangaré et al., "Detection of bacterial pathogens including potential new species in human head lice from Mali," *PLoS One*, vol. 12, no. 9, article e0184621, 2017.
- [8] N. Amanzougaghene, J. Akiana, G. Mongo Ndombe et al., "Head lice of pygmies reveal the presence of relapsing fever *Borreliae* in the Republic of Congo," *PLoS Neglected Tropical Diseases*, vol. 10, no. 12, article e0005142, 2016.
- [9] A. K. Sangaré, C. Socolovschi, M.-M. Olive et al., "Detection of *Bartonella quintana* in African body and head lice," *The American Journal of Tropical Medicine and Hygiene*, vol. 91, no. 2, pp. 294–301, 2014.
- [10] N. Ulutasdemir, F. Eroglu, M. Tanriverdi, E. I. Dagli, and I. S. Koltas, "The epidemic typhus and trench fever are risk for public health due to increased migration in southeast of Turkey," *Acta Tropica*, vol. 178, pp. 115–118, 2018.
- [11] M. Moshki, F. Zamani-Alavijeh, and M. Mojadam, "Correction: efficacy of peer education for adopting preventive behaviors against head lice infestation in female elementary school students: a randomised controlled trial," *PLoS One*, vol. 14, no. 2, article e0212625, 2019.
- [12] D. Awahmukalah, J. Dinga, and J. N. Nchako, "Pediculosis among urban and rural school children in Kumba, Meme division, south-west Cameroon," *Parassitologia*, vol. 30, no. 2-3, pp. 249–256, 1988.
- [13] E. Ebomoyi, "Pediculosis capitis among primary schoolchildren in urban and rural areas of Kwara State, Nigeria," *Journal of School Health*, vol. 58, no. 3, pp. 101–103, 1988.
- [14] R. Speare, J. Govere, and D. Durrheim, "The prevalence of pediculosis in rural South African schoolchildren: research in action," *South African Journal of Science*, vol. 99, no. 1, pp. 21–23, 2003.
- [15] T. A. Abd El Raheem, N. A. El Sherbiny, A. Elgameel, G. A. El-Sayed, N. Moustafa, and S. Shahan, "Epidemiological comparative study of pediculosis capitis among primary school children in Fayoum and Minofiya governorates, Egypt," *Journal of Community Health*, vol. 40, no. 2, pp. 222–226, 2015.
- [16] J. Heukelbach and U. S. Ugbomoiko, "Knowledge, attitudes and practices regarding head lice infestations in rural Nigeria," *The Journal of Infection in Developing Countries*, vol. 5, no. 9, pp. 652–657, 2011.
- [17] G. M. Kirchofer, J. H. Price, and S. K. Telljohann, "Primary grade teachers' knowledge and perceptions of head lice," *Journal of School Health*, vol. 71, no. 9, pp. 448–452, 2001.
- [18] J. H. Price, C. N. Burkhart, C. G. Burkhart, and R. Islam, "School nurses' perceptions of and experiences with head lice," *Journal of School Health*, vol. 69, no. 4, pp. 153–158, 1999.
- [19] E. Sidoti, F. Bonura, G. Paolini, and G. Tringali, "A survey on knowledge and perceptions regarding head lice on a sample of teachers and students in primary schools of north and south of Italy," *Journal of Preventive Medicine and Hygiene*, vol. 50, no. 3, 2009.
- [20] M. Counahan, R. M. Andrews, H. Weld, H. Walsh, and R. Speare, "What parents in Australia know and do about head lice," *Rural and Remote Health*, vol. 7, article 687, p. 3, 2007.
- [21] A. Doulgeraki and M. Valari, "Parental attitudes towards head lice infestation in Greece," *International Journal of Dermatology*, vol. 50, no. 6, pp. 689–692, 2011.
- [22] J. Parison and D. V. Canyon, *Head Lice and the Impact of Knowledge, Attitudes and Practices—a Social Science Overview*, UNI-MED, 2010.
- [23] K. Mumcuoglu, M. Mumcuoglu, M. Danilevich, and L. Gilead, "Knowledge and practices of health professionals regarding head lice," in *Management and control of head lice infestations*, pp. 111–114, UNI-MED Verlag AG, Bremen, 2010.
- [24] B. Olowokure, H. Jenkinson, M. Beaumont, and H. Duggal, "The knowledge of healthcare professionals with regard to the treatment and prevention of head lice," *International Journal of Environmental Health Research*, vol. 13, no. 1, pp. 11–15, 2003.
- [25] Z. Philips, D. Whynes, S. Parnham, R. Slack, and S. Earwicker, "The role of community pharmacists in prescribing medication for the treatment of head lice," *Journal of Public Health*, vol. 23, no. 2, pp. 114–120, 2001.
- [26] Z. G. Shirvani and M. S. Ardestani, "Evaluation of a health education program for head lice infestation in female primary school students in Chabahr City, Iran," *Archives of Iranian Medicine*, vol. 16, no. 1, p. 42, 2013.
- [27] K. Y. Mumcuoglu, L. Gilead, and A. Ingber, "New insights in pediculosis and scabies," *Expert Review of Dermatology*, vol. 4, no. 3, pp. 285–302, 2014.
- [28] R. Pourbaba, M. Moshkbide Haghghi, and P. R. Habibi, "A survey of prevalence of pediculosis among primary school students of Guilan Province in the school year of 2002-3," *Journal of Guilan University of Medical Sciences*, vol. 13, no. 52, pp. 15–24, 2005.
- [29] H. Dagne, A. A. Biya, A. Tirfie, W. W. Yallew, and B. Dagnaw, "Prevalence of pediculosis capitis and associated factors among schoolchildren in Woreta town, northwest Ethiopia," *BMC Research Notes*, vol. 12, no. 1, p. 465, 2019.
- [30] W. N. Arifin, "Introduction to sample size calculation," *Education in Medicine Journal*, vol. 5, no. 2, 2013.
- [31] J. Cohen, *The T Test for Means. Statistical Power Analysis for the Behavioural Sciences*, Lawrence Erlbaum Associates, Hillsdale, NJ, 1988.
- [32] H. Kassiri and E. Esteghali, "Prevalence rate and risk factors of *Pediculus capitis* among primary school children in Iran," *Archives of Pediatric Infectious Diseases*, vol. 4, no. 1, 2015.
- [33] B. Norsa Adah, B. Rusli, I. Aini, and L. Naing, "Is health education necessary to control head lice infestation in primary school children?," *International Medical Journal-Tokyo*, vol. 13, no. 1, p. 19, 2006.
- [34] I. Zareban, M. Abbaszadeh, M. Moodi, H. Mehrjoufard, and H. Ghaffari, *Evaluating a Health-Education Program in Order to Reduce Infection to *Pediculus Humanus Capitis* among Female Elementary Students*, JOURNAL OF BIRJAND UNIVERSITY OF MEDICAL SCIENCES, Scientific information database, 2006.

- [35] N. ul Haq, M. A. Hassali, A. A. Shafie et al., "A cross-sectional assessment of knowledge, attitude and practice among Hepatitis-B patients in Quetta, Pakistan," *BMC Public Health*, vol. 13, no. 1, article 448, 2013.
- [36] A. Singh and B. Purohit, "Knowledge, Attitude and Practice towards Infection Control Measures and its Correlation among Dental Students in Bhopal city, Central India," *International Journal of Infection Control*, vol. 7, no. 1, 2011.
- [37] L. Amelia, C. Anwar, and W. Wardiansah, "Association of sociodemographic, knowledge, attitude and practice with pediculosis capitis," *Bioscientia Medicina: Journal of Biomedicine and Translational Research*, vol. 3, no. 1, pp. 51–63, 2019.
- [38] A. M. Khamaiseh, "Head lice among governmental primary school students in southern Jordan: prevalence and risk factors," *Journal of Global Infectious Diseases*, vol. 10, no. 1, pp. 11–15, 2018.
- [39] M. Al Bashtawy and F. Hasna, *Pediculosis capitis among primary-school children in Mafraq Governorate*, Eastern Mediterranean Health Journal/La Revue de Santé de la Méditerranée orientale, Jordan, 2012.
- [40] M. Davarpanah, D. Mehrabani, F. Khademolhosseini, A. Mokhtari, H. Bakhtiari, and R. Neirami, "The prevalence of *Pediculus capitis* among school children in Fars Province, Southern Iran," *Iranian Journal of Parasitology*, vol. 4, pp. 48–53, 2009.
- [41] M. Gulgun, E. Balci, A. Karaoglu, O. Babacan, and T. Türker, "Pediculosis capitis: prevalence and its associated factors in primary Schoolchildren living in rural and urban areas in Kaiseri, Turkey," *Central European Journal of Public Health*, vol. 21, no. 2, pp. 104–108, 2013.
- [42] A. Buczek, D. Markowska-Gosik, D. Widomska, and I. M. Kawa, "Pediculosis capitis among schoolchildren in urban and rural areas of eastern Poland," *European Journal of Epidemiology*, vol. 19, no. 5, pp. 491–495, 2004.
- [43] S. Willems, H. Lapeere, N. Haedens, I. Pasteels, J.-M. Naeyaert, and J. De Maeseneer, "The importance of socio-economic status and individual characteristics on the prevalence of head lice in schoolchildren," *European Journal of Dermatology*, vol. 15, no. 5, pp. 387–392, 2005.
- [44] H. Lesshafft, A. Baier, H. Guerra, A. Terashima, and H. Feldmeier, "Prevalence and risk factors associated with pediculosis capitis in an impoverished urban community in Lima, Peru," *Journal of Global Infectious Diseases*, vol. 5, no. 4, pp. 138–143, 2013.
- [45] R. B. Nordin, W. A. Che, N. A. Ibrahim, and L. Naing, "Prevalence and associated factors of head lice infestation among primary schoolchildren in Kelantan, Malaysia," *Southeast Asian Journal of Tropical Medicine and Public Health*, vol. 37, no. 3, p. 536, 2006.
- [46] V. Soultana, P. Euthumia, M. Antonios, and R. S. Angeliki, "Prevalence of pediculosis capitis among schoolchildren in Greece and risk factors: a questionnaire survey," *Pediatric Dermatology*, vol. 26, no. 6, pp. 701–705, 2009.