

# Variation in mitochondrial minichromosome composition among *Hoplopleura* lice (Phthiraptera: Hoplopleuridae) from rats

Yi-Tian Fu

Hunan Agricultural University

Yu Nie

Hunan Agricultural University

De-Yong Duan

Hunan Agricultural University

Guo-Hua Liu (✉ [liuguohua5202008@163.com](mailto:liuguohua5202008@163.com))

Hunan Agricultural University <https://orcid.org/0000-0002-8434-899X>

---

## Short report

**Keywords:** Rat louse, Mitochondrial genome, Minichromosome, Phylogenetic analyses

**DOI:** <https://doi.org/10.21203/rs.3.rs-42350/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

---

## Abstract

## Background

The family Hoplopleuridae contains at least 183 species of blood-sucking lice, which widely parasitize both mice and rats. Fragmented mitochondrial (mt) genomes have been reported in two rat lice (*Hoplopleura kitti* and *H. akanezumi*) from this family, but some minichromosomes were unidentified in their mt genomes.

## Methods

We sequenced the mt genome of rat louse *Hoplopleura* sp. with an Illumina HiSeq platform and compared its mt genome organization with *H. kitti* and *H. akanezumi*.

## Results

Fragmented mt genome of the rat louse *Hoplopleura* sp. contains 37 genes which are on 12 circular mt minichromosomes. Each mt minichromosome is 1.8–2.7 kb long, which contains 1–5 genes and one large non-coding region. The gene content and arrangement of three mt minichromosomes of *Hoplopleura* sp. and *H. kitti* are different from that of the three mt minichromosomes of *H. akanezumi*. Phylogenetic analyses based on the deduced amino acid sequences of the eight protein-coding genes showed that the *Hoplopleura* sp. was more closely related to *H. akanezumi* than to *H. kitti*, and then they form a monophyletic group.

## Conclusions

Comparison among the three rat lice revealed variation in the composition of mt minichromosomes within the genus *Hoplopleura*. *Hoplopleura* sp. is the first species from the family Hoplopleuridae for which a complete fragmented mt genome has been sequenced. The new data provides useful genetic markers for studying the population genetics, molecular systematics and phylogenetics of blood-sucking lice.

## Background

Blood-sucking lice are known vectors and transmit various disease agents and cause significant vector-borne diseases in humans, domestic and wild mammals [1]. The family Hoplopleuridae contains at least 183 described species of blood-sucking lice and is currently classified into eight genera [2]. Of the eight genera, *Hoplopleura* Enderlein, 1904 is the most species-rich (165 described species) found on rodents [3]. The *Hoplopleura* spp. is common ectoparasite of both mice and rats, causing pruritus, alopecia, dermal irritation and even anemia.

Metazoan mitochondrial (mt) genomes are usually circular DNA molecules (13–20 kb) with 36–37 genes that contain 12–13 protein-coding genes, two rRNA genes and 22 tRNA genes [4]. However, some parasitic lice have an unusual, fragmented mt genome organization. Fragmentation of the mt genome was first found in the human body louse, *Pediculus humanus corporis* (suborder Anoplura) [5]. Since then, 11 other blood-sucking lice (suborder Anoplura), eight avian feather lice (suborder Ischnocera) and the elephant louse, *Haematomyzus elephantis* (suborder Rhynchophthirina), have been found with fragmented mt genomes [6–15]. To date, the complete mt genomes of 12 blood-sucking lice have been sequenced and deposited in GenBank, but the complete mt genomes have been only reported for two rat lice (*H. kitti* and *H. akanezumi*) from this family Hoplopleuridae [7]. In addition, three genes (*nad1*,

*nad3* and *nad5*) or minichromosomes were unidentified in the mt genomes of two *Hoplopleura* species [7]. Interestingly, gene rearrangement has been reported in fragmented mt genome of two *Hoplopleura* species [7]. Therefore, *Hoplopleura* mt genomes may represent one of the most frequently rearranged/fragmentations mt genomes within the family Hoplopleuridae.

To understand the composition of mt minichromosomes among species of the same genus *Hoplopleura*. We sequenced the complete mt genome of rat louse *Hoplopleura* sp. and compare its mt genome organization with other two *Hoplopleura* species, and to re-construct its phylogenetic relationships within the suborder Anoplura using protein sequences derived from coding genes.

## Methods

### Sample collection and DNA extraction

Adult specimens of *Hoplopleura* sp. were collected from the Edward's long-tailed rats *Leopoldamys edwardsi* in Chongqing, China. The specific identity of the examined wild rats was determined by PCR-based sequencing of the mitochondrial (mt) *cox1* gene using an established method [16]. These rat lice were washed five times in physiological saline solution, identified preliminarily to the genus level (as *Hoplopleura* sp.) based on morphological features [17], and stored in 70% (v/v) ethanol at -20 °C. Whole genomic DNA including nuclear and mt DNA was extracted from 40 single rat lice (25 females and 25 males) using the DNeasy Tissue Kit (Promega Corporation, Madison, USA) according to the manufacturer's recommendations. The identity of these specimens was further confirmed by polymerase chain reaction (PCR) amplification and subsequent sequencing of the mt *cox1* and *rns* genes using primer pairs L6625 (5'-CCGGATCCTTYTGRTTYTTYGGNCAAYCC-3') - H7005 (5'-CCGGATCCACNACRTARTANGTRTCRTG-3') and 12SA (5'-TACTATGTTACGACTTAT-3') - 12SB (5'-AAACTAGGATTAGATACCC-3'), respectively.

### Sequencing and assembling

The purity of the extracted whole genomic DNA was assessed by agarose-gel electrophoresis. The DNA concentration was determined using a Quantus Fluorometer (Invitrogen, UK). A paired-end genomic DNA library (350 bp inserts) was constructed for high throughput sequencing with HiSeq×Ten (Illumina, San Diego, CA, USA) and collected raw reads were exported in the FASTQ format. The raw reads were filtered by removing adaptor reads, redundant reads and 'N'-rich reads. Finally, 2 Gb clean data (250 bp pair-end reads) was produced for this rat louse. Illumina sequence reads were assembled into contigs with Geneious 11.1.5 [18] based on *cox1* and *rns* relatively conserved sequences. The assembly parameters were minimum overlap identity 99% and minimum overlap 100 bp. When the two ends of the contig overlapped, indicating circular organization of the minichromosome. The conserved non-coding region sequences were identified between the *rnsL* and *cox1* minichromosomes and were used as references to align the Illumina sequence dataset. We assembled these minichromosomes individually in full length using the same method stated above for *cox1* and *rns* minichromosome assembly.

### Annotation

Sequences were aligned against the mt minichromosome sequences of rat louse *H. kitti* [7] available using the MAFFT 7.122 software [19] to identify gene boundaries. Protein-coding genes and rRNA genes were identified with BLAST searches of NCBI database. Amino acid sequences of each protein-coding genes were inferred using MEGA 6.0 [20]. tRNA genes were identified using ARWEN [21] and the program tRNAscan-SE [22] with manual adjustment.

### Verification of mt minichromosomes

The size of each mt minichromosome of *Hoplopleura* sp. were verified by PCR using specific primers (Table 1). The forward primer and reverse primer in each pair were next to each other with a small gap in between (10-50 bp). PCR with these primers amplified each circular minichromosome in full length (Fig. 1). To obtain full-length sequences of the non-coding regions of the minichromosomes, these amplicons were also sequenced with high throughput sequencing as described above.

## Phylogenetic analysis

Phylogenetic relationship among representing the blood-sucking lice of suborder Anoplura was performed based on concatenated amino acid sequences (Table 2), using one elephant louse, *H. elephantis* (GenBank accession numbers: KF933032-41) as an outgroup [10]. Eight amino acid sequences (except for *nad1*, *nad2*, *nad3*, *nad4* and *nad5* because these genes were unidentified in some blood-sucking lice) were aligned individually using MAFFT 7.122 and were then concatenated to form a single dataset; ambiguously aligned regions were excluded using Gblocks 0.91b using default parameters [23]. The MtArt + I + G + F was selected as the most appropriate evolutionary model by ProtTest 2.4 based on the Akaike information criterion (AIC) [24]. Phylogenetic analyses were conducted with maximum likelihood (ML) using PhyML 3.0 with a BioNJ starting tree, and tree topology search was set as the best from the nearest neighbor interchange (NNI) and subtree pruning and regrafting (SPR) methods [25]. Bootstrap was calculated using 100 bootstrap replicates. Phylograms were drawn using FigTree v.1.31.

## Results And Discussion

### Identity of the rat louse *Hoplopleura* sp.

Two blood-sucking louse species (*H. kitti* and *Polyplax insulsa*) parasitize in *L. edwardsi* (<http://phthiraptera.info/category/mammal-wilson-reeder/mammals/rodentia/muridae/murinae/leopoldamys/leopoldamys-edwardsi>). The *Hoplopleura* sp. has close morphological and morphometric similarities with *H. kitti* recovered from the same host (*L. edwardsi*). The mt *cox1* and *rns* genes of *Hoplopleura* sp. shared 76% and 77.6% identity with previously published sequences of *H. kitti* (KJ648943) from *Berylmys bowersi* and *H. akanezumii* (KJ648928) from *Apodemus chevrieri* in China and, respectively.

### General features of the mt genome of the rat louse *Hoplopleura* sp.

We sequenced the *Hoplopleura* sp. genome and produced 3 Gb of Illumina short-read sequence data and obtained a total of 6,526,349×2 raw reads from adults of *Hoplopleura* sp.. After quality filtration, 3,937,826×2 clean reads (2 Gb) were generated for assembly of the mt genome. We assembled these sequence-reads into contigs and identified 37 mt genes typical of bilateral animals (Fig. 2; Table 3). These genes are on 12 minichromosomal; each minichromosome is 1.8-2.7 kb in size and consists of a coding region and a non-coding region (NCR) in a circular organization (Table 3). The coding regions have 1-5 genes each and vary in size from 675 bp to 1,760 bp (Table 3). All genes are transcribed in the same direction except for *nad1* gene. The nucleotide sequences of the mt minichromosomes of *Hoplopleura* sp. were deposited in GenBank under accession numbers UM012986 -97.

We sequenced the full-length non-coding regions of all of the 12 mt minichromosomes of the *Hoplopleura* sp., which range from 935 (H-*nad5*-F minichromosome) to 1,305 bp (C-*nad6*-W-L<sub>2</sub> minichromosome) (Table 3). The longest non-coding region of *Hoplopleura* sp. was shorter than the longest non-coding region of other sucking lice known, such as pig lice (2,370 bp) [6] and horse lice (3,276 bp) [13]. As in the human lice [12], rat lice [7] and pig lice [6], each coding region of *Hoplopleura* sp. is flanked by a conserved non-coding AT-rich motif (88 bp,71.6%) upstream and a GC-rich

motif (39 bp, 79.5%) downstream, indicating functional significance of these motifs in the mt genomes of blood-sucking lice.

## Annotation

The boundaries between protein-coding genes of the mt genome of *Hoplopleura* sp. were determined by aligning its sequence and by identifying translation initiation and termination codons with those of *H. kitti* and *H. akanezumii* [7]. *Hoplopleura* sp. mt genome encoded 13 protein-encoding genes. It has four initiation codons (ATT, ATG, TTG, GTG). Among them, both ATT (*nad2*, *nad4L*, *nad5*, *cox3* and *cytb*) and ATG (*nad3*, *nad4*, *nad6*, *atp6* and *atp8*) are the highest frequency of being used as initiation codons. Moreover, TTG (*nad1* and *cox2*) and GTG (*cox1*) are used in the mt genome. This mt genome has three termination codons (TAA, TAG, T). Among them, TAG is the most frequently used with five times altogether, by *cox1*, *nad2*, *nad3*, *nad4L* and *cytb*. TAA with secondary high rate of recurrence (four times) as termination codons, *cox2*, *atp6*, *atp8* and *nad4*, used it in the mt genome of *Hoplopleura* sp.. Furthermore, the genes of *cox3*, *nad1*, *nad5* and *nad6* use T as termination codons. Incomplete termination of protein-coding genes is commonly found in other mt genomes of blood-sucking lice, including *Haematopinus suis* [6], *H. apris* [6], *H. asini* [13], *H. kitti* [7], *P. asiatica* [8], *P. spinulosa* [8], *Pediculus schaeffi* [9], *Microthoracius praelongiceps* [11] and *Pthirus pubis* [12]. In the mt genome of *Hoplopleura* sp., the sizes of the *rrnL* and *rrnS* genes were 1,125 bp and 675 bp, respectively. The 22 tRNA genes ranged from 59 to 71 bp in size. The secondary structures predicted (not shown) were similar to those of *H. kitti* and *H. akanezumii* [7].

## Variation in mt minichromosome composition among three rat lice

The complete mt genome sequences of *Hoplopleura* sp. fragmented into 12 circular minichromosomes. The incomplete mt genomes of *H. kitti* and *H. akanezumii* have identified 11 circular minichromosomes [7]. 11 minichromosomes of the rat louse, *Hoplopleura* sp., have the same gene content and gene arrangement as their counterparts of the rat louse, *H. kitti*. Eight of these minichromosomes of the rat lice, *Hoplopleura* sp. and *H. kitti*, have the same gene content and gene arrangement as their counterparts of the rat louse, *H. akanezumii* [7]. The other two minichromosomes of the rat louse *Hoplopleura* sp., however, are not present in the rat louse *H. akanezumii* [7]. In the *Hoplopleura* sp., one of the minichromosomes has four genes, D-Y-*cox2*-T (Fig. 2). In the *H. akanezumii*, however, this minichromosome has only three genes, D-Y-*cox2*. Similarly, another minichromosome of the *Hoplopleura* sp. has five genes, R-*nad4L*-P-*cox3*-A (Fig. 2). In the *H. akanezumii*, however, this minichromosome has six genes, R-*nad4L*-P-*cox3*-A-T. Interestingly, a chimeric minichromosome has found in the *H. akanezumii* which contains parts of the two rRNA genes, *rrnL* and *rrnS*, which are only 5% (51 bp) and 24% (172 bp) of the full-length *rrnL* and *rrnS*, respectively [7]. However, this case has unidentified in the *H. kitti* and *Hoplopleura* sp..

## Comparative mt genomic analyses of *Hoplopleura* sp. with *H. kitti* and *H. akanezumii*

A comparison of the nucleotide and the amino acid sequences of each protein-encoding gene (except for *nad1*, *nad3* and *nad5*) of the three *Hoplopleura* species is given in Table 4. Pairwise comparisons of the nucleotide and amino acid sequences revealed identities of 50.6-77.2% and 37.5-90.2% among them, respectively. The greatest nucleotide variation was in the *atp8* gene (49.4%), whereas least differences (22.8%) was detected in the *cox1* gene (Table 4). The difference across both concatenated nucleotide and amino acid sequences of the ten protein-coding was 37.5% and 36.8% between *Hoplopleura* sp. and *H. kitti*, 36.7% and 34.7% between *Hoplopleura* sp. and *H. akanezumii*, and 34.6% and 33.4% between *H. kitti* and *H. akanezumii*.

## Phylogenetic relationships

In the present study, phylogenetic analysis of the concatenated amino acid sequence datasets for eight mt protein-coding genes (Fig. 3) showed that the family Hoplopleuridae (*Hoplopleura* sp., *H. kitti* and *H. akanezumi*) clustered to the exclusion of representatives of the families Polyplacidae (*P. asiatica* and *P. spinulosa*), Haematopinidae (*H. apri*, *H. asini* and *H. suis*), Pediculidae (*P. humanus corporis*, *P. humanus capitis* and *P. schaeffi*), Pthiridae (*P. pubis*), and the family Microthoraciidae (*M. praelongiceps*) clustered separately with strong nodal support (Bootstrap = 100). Within the family Hoplopleuridae, *Hoplopleura* sp. and *H. akanezumi* clustered together with moderate support (Bootstrap = 73), to the exclusion of *H. kitti*.

Many studies have indicated that the mt genome sequence is a valuable genetic marker for phylogenetic studies at various taxonomic levels of different organisms, including lice [14,15]. The mt genome sequences of rat louse *Hoplopleura* sp. could promote to reassess the systematic relationships of lice within suborder Anoplura using mt genomic datasets. No species from the other genera (*Ancistroplax*, *Ferrisella*, *Haematopinoidea*, *Paradoxophthirus*, *Pterophthirus*, *Schizophthirus* and *Typhlomyophthirus*) within family Hoplopleuridae was included in our analyses. Therefore, more expanding taxa sampling is necessary for future phylogenetic studies of family Hoplopleuridae using mt genomic dataset.

## Conclusions

Variation in the composition of mt minichromosomes among species of the genus *Hoplopleura*. *Hoplopleura* sp. is the first species from the family Hoplopleuridae for which a complete fragmented mt genome has been sequenced. The new data provides useful genetic markers for studying the population genetics, molecular systematics and phylogenetics of blood-sucking lice.

## Abbreviations

mt: mitochondrial; rDNA:ribosomal DNA; *nad1*:NADH dehydrogenase subunit 1; *nad3*:NADH dehydrogenase subunit 3; *nad5*:NADH dehydrogenase subunit 5; *cox1*:cytochrome c oxidase subunit 1; *rrnS*:small subunit of rRNA; *rrnL*:large subunit of rRNA; tRNA:transfer RNA; *nad2*:NADH dehydrogenase subunit 2; *nad4*:NADH dehydrogenase subunit 4; *nad4L*:NADH dehydrogenase subunit 4L; *cox3*:cytochrome c oxidase subunit 3; *cytb*:cytochrome b; *nad6*:NADH dehydrogenase subunit 6; *atp6*:ATP synthase F0 subunit 6; *atp8*:ATP synthase F0 subunit 8; *cox2*:cytochrome c oxidase subunit 2.

## Declarations

### Acknowledgements

Not applicable.

### Authors' contributions

Y-TF, G-HL conceived and designed the study, and critically revised the manuscript, and Y-TF performed the experiments. Y-TF, G-HL analyzed the data. D-YD and G-HL drafted the manuscript. YN helped in study design, study implementation, and manuscript preparation. All authors read and approved the final manuscript.

### Funding

This study was supported by the Planned Programme of Hunan Province Science and Technology Innovation (Grant no. 2018RS3085) and the Training Programme for Excellent Young Innovators of Changsha (Grant No. KQ1905013).

## Availability of data and materials

The fragmented mitochondrial genome sequences of *Hoplopleura* sp. from the Edward's long-tailed rats have been deposited in the GenBank database under the accession numbers UM012986 -97.

## Ethics approval and consent to participate

All procedures involving animals in the present study were approved and this study was approved by the Animal Ethics Committee of Hunan Agricultural University (No. 43321503).

## Consent for publication

Not applicable.

## Competing interests

The authors declare that they have no competing interests.

## References

1. Tsai YL, Chang CC, Chuang ST, Chomel BB. *Bartonella* species and their ectoparasites: selective host adaptation or strain selection between the vector and the mammalian host? *Comp Immunol Microbiol Infect Dis.* 2011;34:299–314.
2. Kim KC, Ludwig HW. The family classification of the Anoplura. *Syst Entomol.* 1978;3:249–84.
3. Durden LA, Musser GG. The sucking lice (Insecta, Anoplura) of the world: a taxonomic checklist with records of mammalian hosts and geographical distributions. *Bull Amer Mus Nat Hist.* 1994;218:1–90.
4. Boore JL. Animal mitochondrial genomes. *Nucleic Acids Res.* 1999;27:1767–80.
5. Shao R, Kirkness EF, Barker SC. The single mitochondrial chromosome typical of animals has evolved into 18 minichromosomes in the human body louse, *Pediculus humanus*. *Genome Res.* 2009;19:904–12.
6. Jiang H, Barker SC, Shao R. Substantial variation in the extent of mitochondrial genome fragmentation among blood-sucking lice of mammals. *Genome Biol Evol.* 2013;5:1298–308.
7. Dong WG, Song S, Guo XG, Jin DC, Yang QQ, Barker SC, et al. Fragmented mitochondrial genomes are present in both major clades of the blood-sucking lice (suborder Anoplura): evidence from two *Hoplopleura* rodent lice (family Hoplopleuridae). *BMC Genom.* 2014;15:751.
8. Dong WG, Song S, Jin DC, Guo XG, Shao R. Fragmented mitochondrial genomes of the rat lice, *Polyplax asiatica* and *Polyplax spinulosa*: intra-genus variation in fragmentation pattern and a possible link between the extent of fragmentation and the length of life cycle. *BMC Genom.* 2014;15:44.
9. Herd KE, Barker SC, Shao R. The mitochondrial genome of the chimpanzee louse, *Pediculus schaeffi*: insights into the process of mitochondrial genome fragmentation in the blood-sucking lice of great apes. *BMC Genom.* 2015;16:661.
10. Shao R, Barker SC, Li H, Song S, Poudel S, Su Y. Fragmented mitochondrial genomes in two suborders of parasitic lice of eutherian mammals (Anoplura and Rhynchophthirina, Insecta). *Sci Rep.* 2015;5:17389.
11. Shao R, Li H, Barker SC, Song S. The mitochondrial genome of the guanaco louse, *Microthoracius praelongiceps*: insights into the ancestral mitochondrial karyotype of sucking lice (Anoplura, Insecta). *Genome Biol Evol.* 2017;9:431–45.

12. Shao R, Zhu XQ, Barker SC, Herd K. Evolution of extensively fragmented mitochondrial genomes in the lice of humans. *Genome Biol Evol.* 2012;4:1088–101.
13. Song SD, Barker SC, Shao R. Variation in mitochondrial minichromosome composition between blood-sucking lice of the genus *Haematopinus* that infest horses and pigs. *Parasit Vectors.* 2014;7:144.
14. Sweet AD, Johnson KP, Cameron SL. Mitochondrial genomes of *Columbicola* feather lice are highly fragmented, indicating repeated evolution of minicircle-type genomes in parasitic lice. *PeerJ.* 2020;8:e8759.
15. Song F, Li H, Liu GH, Wang W, James P, Colwell DD, et al. Mitochondrial genome fragmentation unites the parasitic lice of eutherian mammals. 2019;68:430 – 40.
16. Robins JH, Hingston M, Matisoo-Smith E, Ross HA. Identifying *Rattus* species using mitochondrial DNA. *Mol Ecol Notes.* 2007;7:717–29.
17. Kim KC, Ludwig HW. The family classification of the Anoplura. *Syst Entomol.* 1978;3:249–84.
18. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, et al. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics.* 2012;28:1647–9.
19. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol.* 2013;30:772–80.
20. Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. MEGA6: Molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol.* 2013;30:2725–9.
21. Laslett D, Canbäck B. ARWEN: a program to detect tRNA genes in metazoan mitochondrial nucleotide sequences. *Bioinformatics.* 2008;24:172–5.
22. Lowe TM, Chan PP. tRNAscan-SE On-line: integrating search and context for analysis of transfer RNA genes. *Nucleic Acids Res.* 2016;44:W54-7.
23. Talavera G, Castresana J. Improvement of phylogenies after removing divergent and ambiguously aligned blocks from protein sequence alignments. *Syst Biol.* 2007;56:564–77.
24. Abascal F, Zardoya R, Posada D. ProtTest: selection of best-fit models of protein evolution. *Bioinformatics.* 2005;21:2104–5.
25. Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst Biol.* 2010;59:307–21.

## Tables

**Table 1** PCR primers used to amplify and sequence the mitochondrial genome of the rat lice, *Hoplopleura* sp.



Primer	Sequence (5' to 3')	Minichromosome
1F	AGCACTTGTTCTGATTCTTCGGTC	<i>I-cox1</i>
1R	TCGTGATACCCCCTGCCAAACTG	<i>I-cox1</i>
2F	CTTTCAAGAGACACAAGGGGTTCA	<i>rrnS</i>
2R	TATTTTCCCAGTCCTACAGAGAGC	<i>rrnS</i>
3F	TGTCCTTGTCCCGAAAGAGAGTGAT	M-L1- <i>rrnL-V</i>
3R	CTATTCCACCCTCCCTGATACAAAA	M-L1- <i>rrnL-V</i>
4F	TGAGTAAGGGGGATACATCACGCTA	Q- <i>nad1-G-nad3</i>
4R	CAGCGAACTCTGCGTATTCCTCCAT	Q- <i>nad1-G-nad3</i>
5F	TAAGGTTATCGGGCATCAGTGGTA	D-Y- <i>cox2-T</i>
5R	AGAGGGGATGGCGAGGACAAAAAG	D-Y- <i>cox2-T</i>
6F	CGCCAACTATCAGAACTTTCCAAC	<i>atp8-atp6-N</i>
6R	TCGTGGATAACAGTCACAAAGATG	<i>atp8-atp6-N</i>
7F	GCATTTACAGTGCTCAGTCTTCGC	<i>nad2</i>
7R	ACAAAGACAAAGGGGGAAACGGGA	<i>nad2</i>
8F	TTAGCGGTAAGCGGGACTGAGGTA	C- <i>nad6-W-L2</i>
8R	AACTCTATTTCCCCGTTTCCCAA	C- <i>nad6-W-L2</i>
9F	GTTCTCTCGGTTTTCCATCCCTCA	R- <i>nad4L-P-cox3-A</i>
9R	TCTATCGCTACCAGAGAGATTGTTA	R- <i>nad4L-P-cox3-A</i>
10F	GGGAAAACCTCCGACAAGGTCACATT	E- <i>cytb-S1-S2</i>
10R	CCTAAGGGATTTGAACTTCCTGTCG	E- <i>cytb-S1-S2</i>
11F	GGTATTGCTAAAGTTTGGAGGTATC	K- <i>nad4</i>
11R	CAGCCAAGAGTATTCTCCCAACAT	K- <i>nad4</i>
12F	GGGGATTACCTCCTTCTCATT	H- <i>nad5-F</i>
12R	AAGCAATGAAGAGCAACAAGGACAC	H- <i>nad5-F</i>

**Table 2** The blood-sucking lice included in the phylogenetic analyses in this study

Species	Hosts	GenBank accession numbers	References
<i>Haematopinus apri</i>	Wild pig	KC814611-19	Jiang et al., 2013
<i>Haematopinus asini</i>	Horse	KF939318, KF939322, KF939324, KF939326, KJ434034-38	Song et al., 2014
<i>Haematopinus suis</i>	Domestic pig	KC814602-10	Jiang et al., 2013
<i>Hoplopleura akanezumi</i>	Rat	KJ648922-32	Dong et al., 2014a
<i>Hoplopleura kitti</i>	Rat	KJ648933-43	Dong et al., 2014a
<i>Microthoradus praelongiceps</i>	Guanacos	KX090378-KX090389	Shao et al., 2017
<i>Pediculus humanus corporis</i>	Human	FJ499473-90	Shao et al., 2009
<i>Pediculus humanus capitis</i>	Human	JX080388-407	Shao et al., 2012
<i>Pediculus schaeffi</i>	Chimpanzee	KC241882-97, KR706168-69	Herd et al., 2015
<i>Pthirus pubis</i>	Human	JQ976018, EU219987-95, HM241895-8	Shao et al., 2012
<i>Polyplax asiatica</i>	Rat	KF647751-61	Dong et al., 2014b
<i>Polyplax spinulosa</i>	Rat	KF647762-72	Dong et al., 2014b
<b><i>Hoplopleura sp.</i></b>	<b>Rat</b>	<b>UM012986 -97</b>	<b>Present study</b>

**Table 3** Mitochondrial minichromosomes of the rat louse *Hoplopleura sp.*, identified by


Illumina sequencing

Minichromosome	Size (bp)	Size of coding region (bp)	Size of non-coding region (bp)	Intergenic region (bp)
<i>I-cox1</i>	2,531	1,549	975	7
<i>rrnS</i>	1,869	675	1,194	0
<i>M-L1-rrnL-V</i>	2,257	1,323	934	0
<i>Q-nad1-G-nad3</i>	2,525	1,445	1,063	17
<i>D-Y-cox2-T</i>	2,087	880	1,129	78
<i>atp8-atp6-N</i>	2,023	896	1,118	9
<i>nad2</i>	2,141	981	1,160	0
<i>C-nad6-W-L2</i>	1,979	673	1,305	1
<i>R-nad4L-P-cox3-A</i>	2,311	1,251	1,057	3
<i>E-cytb-S1-S2</i>	2,417	1,304	1,113	0
<i>K-nad4</i>	2,289	1,313	975	1
<i>H-nad5-F</i>	2,695	1,759	935	1
Total	27,124	14,049	12,958	117

**Table 4** Nucleotide (nt) and/or predicted amino acid (aa) sequence differences in mitochondrial genes among *Hoplopleura* sp. (Hs), *H. kitti* (Hk) and *H. akanezumi* (Ha) upon pairwise comparison


Gene/region	Nt sequence length			Nt difference (%)			Number of aa			aa difference (%)		
	Hs	Hk	Ha	Hs / Hk	Hs / Ha	Hk / Ha	Hs	Hk	Ha	Hs / Hk	Hs / Ha	Hk / Ha
atp6	651	651	654	36.34	36.54	33.49	216	216	217	32.26	33.64	27.65
atp8	174	195	177	47.50	49.44	46.97	57	64	58	62.50	59.02	54.69
nad2	981	990	984	48.44	44.18	43.40	326	329	327	55.15	53.19	54.85
nad4	1,248	1,242	1,254	40.38	40.49	41.21	415	413	417	44.84	45.56	44.84
nad4L	273	273	270	42.34	39.56	42.12	90	90	89	48.89	44.44	50.00
nad6	478	483	474	43.83	44.49	41.74	158	160	157	47.20	50.63	48.75
cox1	1,485	1,530	1,530	29.15	29.26	22.80	494	509	509	17.68	15.32	9.80
cox2	687	681	684	38.24	34.40	33.77	228	226	227	41.30	32.02	31.00
cox3	787	787	789	34.18	34.60	32.57	261	261	262	27.48	29.77	30.92
cytb	1,104	1,102	1,107	34.30	33.97	32.13	367	367	368	29.70	25.27	25.00
rrnS	675	737	690	31.80	25.32	33.73						
rrnL	1,125	1,107	1,131	28.48	27.55	29.86						

## Figures



**Figure 1**

PCR verification of the 12 mt minichromosomes of the rat louse, *Hoplopleura* sp.. M: DL2000 DNA marker; Lane 1-12: I-cox1, rrnS, M-L1-rrnL-V, Q-nad1-G-nad3, D-Y-cox2-T, atp8-atp6-N, nad2, C-nad6-W-L2, R-nad4L-P-cox3-A, E-cytb-S1-S2, K-nad4 and H-nad5-F.



**Figure 2**

The complete mitochondrial genome of rat louse, *Hoplopleura* sp.. Each minichromosome has a coding region and a non-coding region (NCR, in black). The names and transcript orientation of genes are indicated in the coding region and the minichromosomes are placed in alphabetical order of protein-coding genes and rRNA genes. Gene names are all in Abbreviation: *atp6* and *atp8* for ATP synthase subunits 6 and 8; *cob* for cytochrome b; *cox1-3* for cytochrome oxidase subunits 1–3, *nad1-6* and *nad4L* for NADH dehydrogenase subunits 1–6 and 4L; *rrnS* and *rrnL* for small and large subunits of ribosomal RNA. tRNA genes are indicated with their single-letter abbreviations of the corresponding amino acids.




Figure 3

Phylogenetic relationships among 14 species of Anoplura inferred from maximum likelihood of deduced amino acid sequences of eight mitochondrial proteins. One elephant louse, *Haematomyzus elephantis* was used as the outgroup. Bootstrap values were indicated at nodes.